

10/13/00

Jc490 U.S. PRO

10-16-00

A

Jc921 U.S. PRO
10/13/00DIVISION - CONTINUATION - CONTINUATION-IN-PART
APPLICATION TRANSMITTAL FORM

Attorney Docket No.:

A-357C

Anticipated Classification Of This Application:

Class

530

Subclass

Prior Application:

Examiner

S. Gucker

Art Unit

1647

To the Assistant Commissioner for Patents:

This is a request for filing a ☐ continuation ☒ divisional ☐ continuation-in-part application, under 37 CFR 1.53(b), of pending prior application Serial No. 08/535,681 filed on September 28, 19 95.

of Shaw-Fen Sylvia Hufor TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR

For CONTINUATION or DIVISIONAL APPLNS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 1b, below, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

1. ☒ Transmitted herewith are:
- ☒ 82 pages of specification, 8 pages of claim(s) and 1 page of abstract, **totaling** 91 pages.
 - ☒ 9 sheet(s) of drawings.
 - ☒ 2 pages of Oath or Declaration by the applicant(s):
 - ☐ a. Newly executed (original or copy)
 - ☒ b. Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional applns. only)
 - ☒ 22 pages of Sequence Listing
2. ☒ The filing fee is calculated below:

For	Number Filed		Number Extra		Rate		Fee
Total Claims	12	- 20 =	0	x	\$18.00	=	\$ 0.00
Independent Claims	1	- 3 =	0	x	\$80.00	=	0.00
Multiple Dependent Claims	1			+	\$270.00	=	270.00
Basic Fee					\$710.00	=	710.00
Total Filing Fee							\$ 980.00

3. ☒ Please charge Deposit Account No. 01-0519, in the name of Amgen Inc., in the amount of \$ 980.00. An original and one copy are enclosed.
4. ☒ Throughout the prosecution of this application, if any extension of time is necessary, please consider this a request therefor.
5. ☒ The Commissioner is hereby authorized to charge any additional filing fees which may be required by the accompanying application, any additional fees which may be required during pendency of this application as required by 37 CFR 1.16 or 1.17, or credit any overpayment to Deposit Account No. 01-0519 throughout the prosecution of this application.
6. ☒ Cancel in this application original claims 1-30 AND 33-44 of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)

EXPRESS MAIL CERTIFICATE

"Express Mail" mail labeling number: EL36068921USDate of Deposit: October 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

S.L. St. Andrew
Printed Name

S.L. St. Andrew
Signature

7. ☒ Preliminarily, please amend the specification by inserting before the first line the following:

--This application is a ☐ continuation ☒ division of application Serial No. 08/535,681, filed September 28, 1995, which is hereby incorporated by reference.--

8. ☐ Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file. (May only be used if signed by person authorized by § 1.138 and before payment of base issue fee.)

8a. ☒ New formal drawings are enclosed.

9. ☐ Priority of application Serial No. _____ filed on _____ in _____ (country) is claimed under 35 USC 119.

9a. ☐ The certified copy has been filed in prior application Serial No. _____ filed _____

10. ☒ The prior application is assigned of record to Amgen Inc.

11. ☒ A preliminary amendment is enclosed.

12. ☒ Also enclosed Attorney's Statement, Information Disclosure Statement, Modified Form 1449

13. ☐ Other: _____

14. ☒ The power of attorney in the prior application is to:
Ron K. Levy, Registration No.: 31,539; Steven M. Ode, Registration No.: 29,094;
Daniel R. Curry, Registration No.: 32,727


a. ☒ The power appears in the original papers in the prior application.

b. ☐ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.

c. ☒ Address all future communications to
Daniel R. Curry
 at the address below.

Signator: ☐ Assignee of complete interest

☒ Attorney or agent of record


 Daniel R. Curry
 Attorney for Applicant
 Registration No. 32,727
 Phone: (805) 447-8102
 Date: October 13, 2000

Please send all future correspondence to:

U. S. Patent Operations/ DRC
 Dept. 4300, M/S 27-4-A
 AMGEN, INC.
 One Amgen Center Drive
 Thousand Oaks, California 91320-1799

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s): Shaw-Fen Sylvia Hu

Serial No.: To Be Assigned

Group Art Unit No.: Not Yet Assigned

Filed: October 13, 2000

Examiner: Not Yet Assigned

For: TRUNCATED GLIAL CELL LINE-DERIVED
NEUROTROPHIC FACTOR

Docket No.: A-357C

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D. C. 20231

Sir:

Please amend the referenced application as follows:

In the Specification

Page 1, line 2, insert -- This application is a divisional of application Serial No. 08/535,681, filed September 28, 1995 now pending, which is hereby incorporated by reference. --

In the Claims

Please cancel Claims 1-30 and 33-44.

Please amend the following claims:

31. **(Amended)** A method for affecting the survival or function of neurons [of treating Parkinson's Disease] comprising administering [to a patient] a polynucleotide [sequence of Claim 13] encoding a truncated glial cell line-derived neurotrophic factor (GDNF) protein product consisting of an

EXPRESS MAIL CERTIFICATE

"Express Mail" mail labeling number EL360689271US

Date of Deposit: October 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, DC 20231.

S. L. St. Andrew
Printed Name

S. L. St. Andrew
Signature

amino acid sequence

X-(Cys⁴¹-Cys¹³³)-Y

wherein

(Cys⁴¹-Cys¹³³) consists of Cys⁴¹ through Cys¹³³ of SEQ ID NO:2;

Y represents the carboxy terminal group of Cys¹³³, a carboxy-terminus amino acid residue of Ile¹³⁴, or a substituted amino acid residue, and

X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G

RG

NRG

KNRG (SEQ ID NO:3)

GKNRG (SEQ ID NO:4)

RGKNRG (SEQ ID NO:5)

QRGKNRG (SEQ ID NO:6)

GORGKNRG (SEQ ID NO:7)

RGORGKNRG (SEQ ID NO:8)

RRGORGKNRG (SEQ ID NO:9)

G RRGORGKNRG (SEQ ID NO:10)

KG RRGORGKNRG (SEQ ID NO:11)

GKG RRGORGKNRG (SEQ ID NO:12)

RGKG RRGORGKNRG (SEQ ID NO:13)

SRGKG RRGORGKNRG (SEQ ID NO:14)

NSRGKG RRGORGKNRG (SEQ ID NO:15)

ENSRGKG RRGORGKNRG (SEQ ID NO:16)

PENSRGKG RRGORGKNRG (SEQ ID NO:17)

NPENSRGKG RRGORGKNRG (SEQ ID NO:18)

ANPENSRGKG RRGORGKNRG (SEQ ID NO:19)

A ANPENSRGKG RRGORGKNRG (SEQ ID NO:20)

AA ANPENSRGKG RRGORGKNRG (SEQ ID NO:21)

AAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:22)

OAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:23)

ROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:24)

NRQAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:25)

RNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:26)
ERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:27)
RERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:28)
RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:29)
P RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:30)
LP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:31)
VLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:32)
AVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:33)
MAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:34)
QMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:35)
KOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:36)
DKOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:37) and
PDKOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:38)

or a substitution or deletion variant of X, wherein said variant is in excess of 70% identical to an amino acid sequence of X as set forth above when four gaps in a length of 100 amino acids may be introduced to assist in that alignment, to provide *in vivo* production of said truncated GDNF protein.

32. (Amended) A method according to claim 31 [of treating Parkinson's Disease] comprising implanting in a patient a cell transformed with said [a] polynucleotide [sequence of Claim 13] to provide *in vivo* production of said truncated GDNF protein.

Please add the following new claims:

-- 45. A method according to Claim 31 or 32, wherein X is selected from the group consisting of SEQ ID NO: 3, 7, 8, 14, 17 and 18.


46. A method according to Claim 31 or 32, wherein X is G, RG or NRG.

47. A method according to Claim 31 or 32, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:42.

48. A method according to Claim 31 or 32, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:44.

49. A method according to Claim 31 or 32, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:46. --

Respectfully submitted,



Daniel R. Curry
Attorney for Applicant
Registration No.: 32,727
Phone: (805) 447-8102
Date: October 13, 2000

Please send all future correspondence to:

U.S. Patent Operations/ DRC
Dept. 4300, M/S 27-4-A
AMGEN INC.
One Amgen Center Drive
Thousand Oaks, California 91320-1799

TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTORField of the Invention

In general, the present invention relates to proteins, referred to herein as glial cell line-derived neurotrophic factors (also referred to as glial derived neurotrophic factor or GDNF), that are characterized by the ability to promote dopamine uptake by dopaminergic neurons and support the survival of the neurons that die in Parkinson's Disease. The present invention more specifically relates to a novel truncated GDNF proteins.

Background of the Invention

Neurotrophic factors are proteins, found in the nervous system or in non-nerve tissues innervated by the nervous system, whose function is to promote the survival and maintain the phenotypic differentiation of nerve and/or glial cells (Varon et al., *Ann. Rev. Neuroscience* 1:327, 1979; Thoenen et al., *Science* 229:238, 1985). Because of this physiological role, neurotrophic factors are useful in treating the degeneration of nerve cells and the loss of differentiated function that occurs in a variety of neurodegenerative diseases.

In order for a particular neurotrophic factor to be potentially useful in treating nerve damage, the class or classes of damaged nerve cells must be responsive to the factor. Different neurotrophic factors typically affect distinctly different classes of nerve cells. Therefore, it is advantageous to have on hand a variety of different neurotrophic factors to treat each of the classes of damaged neurons that may occur with different forms of disease or injury.

Neurotrophic factors can protect responsive neurons against a variety of unrelated insults. For example, nerve growth factor (NGF) will rescue a significant portion of sensory neurons from death caused by cutting their axonal processes (Rich et al., *J. Neurocytol* 16:261, 1987; Otto et al., *J. Neurosci.* 83:156, 1987), from ontogenetic death during embryonic development (Hamburger et al., *J. Neurosci.* 4:767, 1984), and from damage caused by the administration of taxol or cisplatin (Apfel et al., *Ann. Neurol.* 29: 87, 1991). This apparent generality of protection has led to the concept that if a neurotrophic factor protects responsive neurons against experimental damage, it may be useful in treating diseases that involve damage to those neurons in patients, even though the etiology may be unknown.

"Express Mail" mail labeling number: TB 530269575 US	Date of Deposit: September 28, 1995
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner of Patents, Washington, D.C. 20231. (Typed Name Below)	
(typed name) Ronda G. Spanr	<i>Ronda G. Spanr</i>

A given neurotrophic factor, in addition to having the correct neuronal specificity, must be available in sufficient quantity to be used as a pharmaceutical treatment. Since neurotrophic factors are typically present in small amounts in tissues (e.g., Hofer and Barde *Nature* 331:261, 1988; Lin et al., *Science* 246:1023, 1989), it would be inconvenient to prepare pharmaceutical quantities of neurotrophic factors directly from animal tissues. As an alternative, it is desirable to use a recombinant expression system to produce the desired protein.

Lin et al. previously described a method for screening biological samples for neurotrophic activity on the embryonic precursors of the substantia nigra dopaminergic neurons (see U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254); the disclosures of which are hereby incorporated by reference). This bioassay is useful in identifying neurotrophic factors which may be used in treating Parkinson's disease (Friedman et al., *Neuro. Sci. Lett.* 79:65-72, 1987) as the disease is characterized by the degeneration of dopaminergic neurons in the midbrain that innervate the striatum.

Lin et al. further described the characterization of a new neurotrophic factor that was purified from one such source, the conditioned culture medium from a glioblastoma cell line, B49 (Schubert et al., *Nature* 249:224-27, 1974). The conditioned medium from this cell line was previously reported to contain dopaminergic neurotrophic activity (Bohn et al., *Soc. Neurosci. Abs.* 15:277, 1989). Prior to the disclosure of Lin et al., glial cell line-derived neurotrophic factor (GDNF) had not been identified as a discrete biologically active substance or isolated as a substantially pure protein. In addition, Lin et al. described processes for cloning human genes encoding GDNF, the nucleic acid sequence of the human genes that encode GDNF and the amino acid sequences of the GDNF protein. The GDNF gene was subcloned into an expression vector, and the vector was used to express biologically active GDNF. The GDNF protein is a homodimer composed of two 134 amino acid, 22 kDa, subunits joined by disulfide bond. The description further included the use of GDNF for preventing and treating nerve damage and nerve related diseases such as Parkinson's disease.

GDNF therapy is helpful in the treatment of nerve damage caused by conditions that compromise the survival and/or proper function of one or more types of nerve cells. Such nerve damage may occur from a wide variety of

different causes. Nerve damage may occur to one or more types of nerve cells by:

(1) physical injury, which causes the degeneration of the axonal processes and/or nerve cell bodies near the site of injury; (2) temporary or permanent cessation of blood flow to parts of the nervous system, as in stroke;

(3) intentional or accidental exposure to neurotoxins, such as chemotherapeutic agents (e.g., cisplatinum) for the treatment of cancer or dideoxycytidine (ddC) for the treatment of AIDS; (4) chronic metabolic diseases, such as diabetes or renal dysfunction; or (5) neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease, and Amyotrophic Lateral Sclerosis (ALS), which result from the degeneration of specific neuronal populations.

GDNF therapy could be particularly helpful in the treatment of neurodegenerative conditions involving the degeneration of the dopaminergic neurons of the substantia nigra, such as Parkinson's disease. The only current treatments for Parkinson's disease are palliative, aiming at increasing dopamine levels in the striatum. The expected impact of GDNF therapy is not simply to produce an increase in the dopaminergic neurotransmission at the dopaminergic nerve terminals in the striatum (which will result in a relief of the symptoms), but also to slow down, or even stop, the progression of the degenerative processes and to repair the damaged nigrostriatal pathway and restore its function. GDNF may also be used in treating other forms of damage to or improper function of dopaminergic nerve cells in human patients. Such damage or malfunction may occur in schizophrenia and other forms of psychosis. The only current treatments for such conditions are symptomatic and require drugs which act upon dopamine receptors or dopamine uptake sites, consistent with the view that the improper functioning of the dopaminergic neurons which innervate these receptor-bearing neuronal populations may be involved in the disease process.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides novel truncated glial cell line-derived neurotrophic factor (GDNF) protein products. In one embodiment, truncated GDNF proteins are produced by recombinant genetic engineering techniques. In an alternative embodiment, the truncated GDNF proteins are synthesized by chemical techniques, or a combination of the recombinant and chemical techniques.

The truncated GDNF protein products of the present invention include the proteins represented by the amino acid sequence X-[Cys⁴¹-Cys¹³³]-Y. The amino acid residue numbering scheme of Figure 1 (SEQ ID NO:2) is used to facilitate comparison to the mature GDNF protein. [Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO:2). Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴. X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

10

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GQRGKNRG (SEQ ID NO:7)
RQRGKNRG (SEQ ID NO:8)
RRQRGKNRG (SEQ ID NO:9)
G RRGQRGKNRG (SEQ ID NO:10)
KG RRGQRGKNRG (SEQ ID NO:11)
GKG RRGQRGKNRG (SEQ ID NO:12)
RGKG RRGQRGKNRG (SEQ ID NO:13)
SRGKG RRGQRGKNRG (SEQ ID NO:14)
NSRGKG RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG RRGQRGKNRG (SEQ ID NO:17)
NPENSRGKG RRGQRGKNRG (SEQ ID NO:18)
ANPENSRGKG RRGQRGKNRG (SEQ ID NO:19)
A ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)
AA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:21)
AAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:22)
QAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:23)
RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24)
NRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:25)

	RNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:26)
	ERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:27)
	RERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:28)
	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:29)
P	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:30)
LP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:31)
VLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:32)
AVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:33)
MAVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:34)
QMAVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:35)
KQMAVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:36)
DKQMAVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:37)
PDKQMAVLP	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:38)

It is contemplated that such truncated GDNF protein products would include truncated GDNF protein having the amino acid sequence as represented by X-[Cys⁴¹-Cys¹³³]-Y and variants and derivatives thereof. Thus, the truncated

- 5 GDNF protein products of the present invention also include addition, substitution and internal deletion variants and derivatives of the amino acid sequences represented by X-[Cys⁴¹-Cys¹³³]-Y. The truncated GDNF protein products further include methionylated or nonmethionylated forms as well as glycosylated or non-glycosylated forms of truncated GDNF protein.
- 10 Exemplary truncated GDNF proteins of the present invention include [Arg¹⁶-Ile¹³⁴], [Asn²²-Ile¹³⁴], [Pro²³-Ile¹³⁴], [Ser²⁶-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴], [Lys³⁷-Ile¹³⁴] and [Asn³⁸-Ile¹³⁴] truncated GDNF proteins, either methionylated or nonmethionylated, and variants and derivatives thereof. Presently preferred truncated GDNF proteins
- 15 of the present invention include [Lys³⁷-Ile¹³⁴] and [Asn³⁸-Ile¹³⁴] truncated GDNF proteins, either methionylated or nonmethionylated, and variants and derivatives thereof. Exemplary substitution variants are the [Asn²²ΔSer²²-Ile¹³⁴] and [Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] truncated GDNF proteins. An exemplary addition variant is the Ser-[Pro²³-Ile¹³⁴] truncated GDNF protein.
- 20 In another aspect of the present invention, the truncated GDNF proteins may be made in glycosylated or non-glycosylated forms. Derivatives of truncated GDNF protein typically involve attaching the truncated GDNF protein to a water soluble polymer. For example, the truncated GDNF protein may be conjugated to

one or more polyethylene glycol molecules to decrease the precipitation of the truncated GDNF protein product in an aqueous environment.

Yet another aspect of the present invention includes the various polynucleotides encoding truncated GDNF proteins. These nucleic acid sequences are generally used in the expression of truncated GDNF in a eukaryotic or prokaryotic host cell, wherein the expression product or a derivative thereof is characterized by the ability to increase dopamine uptake by dopaminergic cells. The polynucleotides may also be used in cell therapy or gene therapy applications. Suitable nucleic acid sequences include those specifically depicted in the Figures as well as additional degenerate sequences and naturally occurring allelic variations.

A further aspect of the present invention involves vectors containing the polynucleotides encoding truncated GDNF proteins operatively linked to amplification and/or expression control sequences. Both prokaryotic and eukaryotic host cells may be stably transformed or transfected with such vectors to express the truncated glial derived neurotrophic factor. The present invention further includes the recombinant production of a truncated GDNF protein wherein such transformed or transfected host cells are grown in a suitable nutrient medium, and the truncated GDNF expressed by the cells is, optionally, isolated from the host cells and/or the nutrient medium. The present invention further includes the use of polynucleotides encoding truncated GDNF and vectors containing such polynucleotides in gene therapy or cell therapy.

In another aspect, the present invention involves a recombinantly produced GDNF composition containing a mixture of a mature GDNF protein and one or more truncated GDNF proteins derived therefrom, wherein the mature GDNF protein has a molecular weight of approximately 44 kDa, and wherein the truncated GDNF protein has a molecular weight of approximately 36 to 40 kDa. The GDNF composition may contain at least two truncated GDNF species wherein a first species has a molecular weight of approximately 36 kDa and a second species has a molecular weight of approximately 40 kDa. The truncated GDNF species having a molecular weight of approximately 40 kDa is a heterodimer of a GDNF monomer having a molecular weight of approximately 22 kDa and a truncated GDNF monomer having a molecular weight of approximately 18 kDa. It is also contemplated that one or more of the truncated GDNF species may be isolated from such a mixture for therapeutic use.

Another aspect of the present invention includes pharmaceutical compositions containing truncated GDNF protein product. Typically, the truncated GDNF protein product is formulated in association with a pharmaceutically acceptable vehicle. A variety of other formulation materials may be used to facilitate manufacture, storage, handling, delivery and/or efficacy. In another aspect of the present invention, truncated GDNF protein products increase dopamine uptake and survival of dopaminergic neurons. Thus, the truncated GDNF protein products are particularly suitable for the treatment of damage to the nervous system caused by injury or disease, such as Parkinson's Disease.

Additional aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following description, which details the practice of the present invention.

BRIEF DESCRIPTION OF THE FIGURES

Numerous features and advantages of the present invention will become apparent upon review of the figures, wherein:

Figure 1 depicts a nucleic acid sequence (SEQ ID NO:1) encoding mature human glial cell line-derived neurotrophic factor (hGDNF). Also depicted is the amino acid sequence (SEQ ID NO:2) of the mature human GDNF protein.

Figure 2 depicts a diagram of a plasmid construction made for the expression of recombinant truncated GDNF proteins.

Figure 3 depicts a restriction map of an alternative nucleic acid sequence (SEQ ID NO:39) encoding GDNF and truncated GDNF polynucleotides.

Figure 4 depicts a restriction map of yet another nucleic acid sequence (SEQ ID NO:40) encoding GDNF and truncated GDNF polynucleotides.

Figure 5 depicts a nucleic acid sequence (SEQ ID NO:41) encoding [Pro²³-Lys³⁷ Δ Asn³⁷-Ile¹³⁴] truncated GDNF protein substitution variant

(SEQ ID NO:42). This protein may also be described as a Met-Ser-[Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] truncated GDNF protein addition/substitution variant.

Figure 6 depicts a nucleic acid sequence (SEQ ID NO:43) encoding an [Arg³²-Ile¹³⁴] truncated GDNF protein (SEQ ID NO:44).

Figure 7 depicts a nucleic acid sequence (SEQ ID NO:45) encoding a [Gly³³-Ile¹³⁴] truncated GDNF protein (SEQ ID NO:46).

Figure 8 depicts the amino acid sequence of mature hGDNF (SEQ ID NO:47) in comparison to several exemplary truncated GDNF proteins: Met-[Arg³²-Ile¹³⁴] (SEQ ID NO:48), Met-[Gly³³-Ile¹³⁴] (SEQ ID NO:49) and Met-Ser-[Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] (SEQ ID NO:50).

DETAILED DESCRIPTION OF THE INVENTION

Human glial cell line-derived neurotrophic factor (hGDNF) is synthesized as a precursor that is processed and secreted as a mature protein of 134 amino acids. It was previously determined that mature human GDNF has the amino acid sequence depicted in Figure 1 (SEQ ID NO:2).

The present invention is based on the unexpected discovery that the mature GDNF protein may be reduced in size (also referred to herein as a "clipped" or "truncated" protein or truncated GDNF protein) yet retain its biological activity. The clipped protein was first discovered during the recombinant production of GDNF in Chinese hamster ovary (CHO) cells. In brief, the recombinant human GDNF (rhGDNF) was prepared as follows. A nucleic acid sequence encoding the entire open reading frame of the mature human GDNF protein was cloned into an expression plasmid. The nucleic acid sequence was confirmed to be correct (by DNA sequencing as the equivalent of the hGDNF sequence in GeneBank) and was translated to an amino acid sequence identical to the published sequence for mature human GDNF (Lin et al., *Science* 260, 1130-1132, 1993). The plasmid DNA was linearized and transfected into dihydrofolate reductase-deficient CHO cells (CHO^{d-} cells) using the calcium phosphate precipitation method. Transfected cells were cultured in a selective

medium, and those colonies that survived the selection process were chosen for individual analysis of hGDNF expression.

Serum-free conditioned media from the individual clones were collected and subjected to Western blot analysis using antisera specific for hGDNF. The antisera involved rabbit polyclonal antibodies elicited from rabbits immunized with recombinant hGDNF expressed in *Escherichia coli*. Under reducing conditions, the hGDNF that was present in these samples was resolved into two major bands having apparent molecular weights of approximately 22 kDa and 18 kDa. Each band consisted of a closely spaced doublet of approximately 22 + 22.5 kDa and 18 + 17.5 kDa, respectively (for simplicity, these doublets will be referred as the 22 kDa and 18 kDa bands or species).

GDNF had previously been reported to exist as a disulfide-bonded homodimer composed of two identical subunits of the mature GDNF protein having a molecular weight of approximately 20 to 22 kDa. When GDNF was analyzed under nonreducing conditions, it was reported that a broad band of 32 to 42 kDa (Lin et al., *Science* 260, 1130-1132, 1993) or 33 to 45 kDa (Lin et al., *J. Neurochem.* 63(2), 758-768, 1994) had been identified. The existence of the range was interpreted as being due to the heterogeneity of glycosylation on the mature monomers and was further substantiated with de-glycosylation experiments.

While the present 22 kDa band corresponds to the mature GDNF protein reported in the literature, the 18 kDa band has not previously been reported. The relative amounts of the 22 kDa and the 18 kDa protein varied in samples collected from individual clones. In addition, it was found that multiple harvests from the same clone showed a variable ratio of the two bands. Moreover, it was found that storage of the CHO-expressed GDNF protein frequently led to an increase in the presence of the 18 kDa band with a concurrent decrease of the 22 kDa band.

When conditioned medium from the transformed CHO^d cells was analyzed under nonreducing conditions by Western blots, three well-resolved bands with apparent molecular weights of 36, 40 and 44 kDa were observed. This finding was also in contrast to previous reports. The relative intensity of these bands was variable, but they correlated well with the ratio of the 22 and 18 kDa monomer bands present in each of the samples. Upon further analysis with monoclonal antisera, it was determined that the three bands in the nonreducing gel corresponded to three possible dimers composed of the two monomers. The

largest 44 kDa protein is a dimer of two 22 kDa mature GDNF proteins as previously reported. The intermediate 40 kDa protein consists of a dimer in which one mature protein has been reduced in molecular weight to an 18 kDa form. The smallest 36 kDa dimer appears to contain two 18 kDa proteins, i.e., both 22 kDa forms have been reduced in molecular weight. This data demonstrated for the first time not only the presence of a novel form of GDNF monomer but also the presence of the clipped GDNF protein in the dimeric configuration. It was also found that, when stored, the monomer composition of the samples shifted towards that of the clipped form and the corresponding dimer species, i.e., the amount of the 36 kDa protein was seen to increase.

Studies were then performed to identify which part of the protein was being eliminated or changed to cause of the reduction in molecular weight in comparison to that of the previously reported mature GDNF protein. It was first determined that the reduction in molecular weight was not due to changes in glycosylation.

GDNF contains two potential N-linked glycosylation sites and has been reported to be glycosylated. The clipped protein, however, is not simply the nonglycosylated or underglycosylated form of mature GDNF. This was demonstrated in deglycosylation experiments wherein samples were treated with N-glycanase, O-glycanase and neuraminidase. On reducing gels, the 18 kDa protein was reduced to a 13.5 kDa band by N-glycanase digestion indicating the presence of an equivalency of 4.5 kDa of N-linked sugar. Treatment with neuraminidase and O-glycanase caused the 18 kDa band to shift slightly to 17 kDa. This indicated the presence of O-linked sugars on the protein. The mature 22 kDa band has been reported to be glycosylated and was also reduced to 18 kDa (i.e., also by 4.5 kDa) by N-glycanase. This was further confirmed through the use of a monoclonal antibody which is specific for the 22 kDa band on the gel. The glycanase digestion pattern of the nonreduced dimer was more complicated, but was interpretable and consistent with the initial assignment of the three forms.

As a result, the 4.5 kDa reduction in molecular weight of the protein was then viewed as resulting from the deletion of approximately 30-35 amino acid residues rather than from changes in glycosylation. The deletion was expected to most likely occur at the amino-terminus of the mature GDNF protein for the following reasons. Mature GDNF contains a total of seven cysteines. If the deletion were from the carboxyl terminus, 2 to 4 of the seven cysteines would be lost, and

this would likely result in an inactive protein. However, when a test sample consisting of predominantly the clipped form was subjected to a bioassay to measure its dopaminergic neuron neurotrophic activity, the sample demonstrated comparable activity to a sample which contained proportionally more of the mature form of GDNF.

The site of cleavage was then determined via amino acid sequence analysis of the purified protein. Samples were sequenced, according to manufacturer's instructions, using an Applied Biosystems 494A protein sequencer for ten cycles. While amino acid sequence analysis techniques and procedures are well known to those skilled in the art, further descriptions of the sequencing of proteins are provided in Fausset et al., *Electrophoresis* 12:22-27, 1991 and United States Patent Application No. 576.316 filed August 24, 1990 (European Patent Application No. 90310899, Publication No. EP 423 980, filed October 4, 1990, entitled "Stem Cell Factor") the disclosures of which are hereby incorporated by reference. Upon analysis, it was determined that the amino terminus of the clipped protein was "RGQRGK" or Arg-Gly-Gln-Arg-Gly-Lys. Therefore, the first 31 amino acids had been removed from the mature protein in the conditioned media. The remaining amino acid sequence of the clipped protein, beginning with amino acid Arg³², was otherwise consistent with that of the mature GDNF amino acid sequence depicted in Figure 1 (SEQ ID NO:2).

The [Arg³²-Ile¹³⁴] truncated GDNF protein was found to be active, on a qualitative basis, in a dopaminergic neuron assay. The dopaminergic neurotrophic activity assay is used to identify neurotrophic factors that may be beneficial in treating Parkinson's disease. The assay is based on a previously described assay (Friedman et al., *Neuro. Sci. Lett.* 79:65-72, 1987, the disclosure of which is hereby incorporated by reference) and may include modifications as described in Lin et al. (see U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254)). A detailed description of the assay is provided in Example 5, below.

A subsequent purification procedure, followed by amino acid sequencing, led to the discovery of another protein from which the first 36 amino acid residues had been removed from the N-terminus of mature GDNF: a [Lys³⁷-Ile¹³⁴] truncated GDNF protein, with an N-terminal sequence of KNRG(C)VL--. Again, the remaining amino acid residues of the clipped protein were otherwise

consistent with those of the mature human GDNF amino acid sequence. The [Lys³⁷-Ile¹³⁴] truncated GDNF protein was also analyzed in the dopamine uptake bioassay. This truncated GDNF protein was found to be active with an ED50 of about 50 pg/ml, similar to that of purified recombinant *E. coli*-expressed, mature GDNF.

It was further discovered that bacterially expressed mature GDNF could be changed to a truncated form. Mature GDNF, expressed in transformed *E. coli* (as described in Lin et al., U.S. Patent Application No. 08/182,183, *supra*), was incubated with CHO-derived conditioned media. Recombinant *E. coli* GDNF has an apparent molecular weight of 17 kDa on reducing gel. When the material was mixed with CHO cell conditioned media and incubated for five days at 4°C, the protein was clipped completely to 12.5 kDa. This cleavage was less complete with one hour or 24 hours of incubation, suggesting a time-dependent process under such conditions. It was also found that simply incubating recombinant *E. coli* GDNF overnight with media containing 0.1% fetal bovine serum did not generate the clipped form. Thus, the presence of live cells in the culture seems to be necessary for the clipping process to occur. It is possible, therefore, that the clipping event may also occur *in vivo* within certain tissues.

In addition, it was found that derivatives of mature *E. coli*-expressed hGDNF, such as pegylated GDNF (also described in Lin et al., U.S. Patent Application No. 08/182,183, *supra*) may be processed to a truncated form in the presence of CHO-derived conditioned media. Mature GDNF may be pegylated at the amino terminus in order to enhance its clearance time in circulation. Pegylation increases the size of the protein, and the modified mature GDNF migrates at about 45 kDa under reduced conditions. As with the non-pegylated mature form, the incubation of pegylated *E. coli* GDNF with CHO cell (untransfected) conditioned media generated a 12.5 kDa band. In both cases, the 12.5 kDa species was present as a disulfide-bonded dimer as shown on non-reducing gels. The generation of this clipped form from the N-terminally pegylated mature protein further demonstrated that the clipping event occurred at the N-terminus of the protein since the pegylated residue was lost during the clipping process.

Based on these findings and because the clipping event may also occur *in vivo*, a truncated form of the GDNF protein may be the ultimate naturally processed form of hGDNF under physiological conditions. Therefore, it was considered advantageous to produce a truncated GDNF protein, or derivative

thereof, for therapeutic use. For example, a directly expressed or synthesized truncated GDNF protein, such as the [Arg³²-Ile¹³⁴] truncated GDNF protein, would be expected to be resistant to the above-described proteolytic activity. Moreover, if it was desired to produce a truncated GDNF derivative, such as a

5 pegylated [Arg³²-Ile¹³⁴] truncated GDNF protein, the resulting derivative would be expected to have the advantage of not being susceptible to the specific clipping which was observed with the mature GDNF derivative.


Additional advantages can also be expected of truncated GDNF protein products. First, the pI of a truncated protein, such as the [Arg³²-Ile¹³⁴]

10 truncated GDNF protein, will be reduced from about 10 to about 8.0-8.5. This makes the protein significantly less basic which could in turn provide beneficial effects including better receptor binding and decreased cytotoxicity at the site of administration, such as an intrathecal injection site. Second, within the first 26 amino acids of the mature GDNF amino acid sequence are two deamidation sites:

15 Arg-Asn-Arg (amino acids 14-16) and Glu-Asn-Ser (amino acids 24-26). The absence of one or both of these sites in a truncated GDNF protein is expected to increase the stability of the protein.

Truncated GDNF Protein Products

20 In a basic embodiment, the truncated GDNF proteins of the present invention may be represented by the following amino acid sequence wherein the amino acid residue numbering scheme of Figure 1 is used to facilitate comparison to the mature GDNF protein:



25 wherein

[Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO:2);

Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

30 X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
RG
NRG
KNG (SEQ ID NO:3)

GKNRG (SEQ ID NO:4)
 RGKNRG (SEQ ID NO:5)
 QRGKNRG (SEQ ID NO:6)
 GQRGKNRG (SEQ ID NO:7)
 RGQRGKNRG (SEQ ID NO:8)
 RRGQRGKNRG (SEQ ID NO:9)
 G RRGQRGKNRG (SEQ ID NO:10)
 KG RRGQRGKNRG (SEQ ID NO:11)
 GKG RRGQRGKNRG (SEQ ID NO:12)
 RGKG RRGQRGKNRG (SEQ ID NO:13)
 SRGKG RRGQRGKNRG (SEQ ID NO:14)
 NSRGKG RRGQRGKNRG (SEQ ID NO:15)
 ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
 PENSRGKG RRGQRGKNRG (SEQ ID NO:17)
 NPENSRGKG RRGQRGKNRG (SEQ ID NO:18)
 ANPENSRGKG RRGQRGKNRG (SEQ ID NO:19)
 A ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)
 AA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:21)
 AAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:22)
 QAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:23)
 RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24)
 NRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:25)
 RNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:26)
 ERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:27)
 RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:28)
 P RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:30)
 LP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:31)
 VLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:32)
 AVL P RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:33)
 MAVLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:34)
 QMAVLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:35)
 KQMAVLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:36)
 DKQMAVLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:37)
 PDKQMAVLP RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:38)

As used herein, the term "truncated GDNF protein product" includes biologically active synthetic or recombinant truncated GDNF proteins, truncated GDNF proteins produced from mature GDNF, biologically active truncated GDNF variants (including insertion, substitution and deletion variants), and
5 chemically modified derivatives thereof. Also included are truncated GDNF proteins that are substantially homologous to the human GDNF protein having the amino acid sequence set forth in SEQ ID NO:2.

The term "biologically active" as used herein means that the truncated GDNF protein demonstrates similar neurotrophic properties, but not necessarily
10 all of the same properties, and not necessarily to the same degree, as the GDNF protein having the amino acid sequence set forth in SEQ ID NO:2. The selection of the particular neurotrophic properties of interest depends upon the use for which the truncated GDNF protein product is being administered. The truncated GDNF protein products are biologically active and demonstrate dopaminergic
15 neuron survival characteristics similar to that demonstrated by mature GDNF protein using the evaluation of dopamine uptake and tyrosine hydroxylase (TH) expression as an exemplary bioassay, as discussed in the examples, below.

The term "substantially homologous", as used herein, means a degree of homology to the human GDNF having the amino acid sequence set forth in SEQ ID
20 NO:2 that is preferably in excess of 70%, most preferably in excess of 80%, and even more preferably in excess of 90% or even 95%. The percentage of homology as described herein is calculated as the percentage of amino acid residues found in the smaller of the two sequences which align with identical amino acid residues in the sequence being compared when four gaps in a length of
25 100 amino acids may be introduced to assist in that alignment as set forth by Dayhoff, in *Atlas of Protein Sequence and Structure* Vol. 5, p. 124 (1972), National Biochemical Research Foundation, Washington, D.C., the disclosure of which is hereby incorporated by reference. Also included as substantially
30 homologous is any truncated GDNF protein which may be isolated by virtue of cross-reactivity with antibodies to the GDNF of SEQ ID NO:2 or whose genes may be isolated through hybridization with the gene or with segments of the gene encoding the GDNF of SEQ ID NO:1.

As will be apparent to those skilled in the art upon reading the present description, substantially homologous proteins will involve one or more
35 deletions from, or additions or substitutions to, the amino acid residues of the truncated GDNF protein represented by X-[Cys⁴¹-Cys¹³³]-Y. The production

of such variants is described in further detail below. It will be further appreciated that because the present invention clearly addresses "truncated" GDNF proteins, the amino-terminus addition variants are contemplated as including the addition of a methionine residue, or non-GDNF amino acid residue or sequence, but do not include the addition of an amino acid residue(s) that would result in the reconstruction of the mature GDNF protein. Truncated GDNF proteins based upon naturally occurring allelic mutants or variants are also within the scope of the present invention. The production of variant truncated GDNF protein is described in further detail below.

Lin et al. (U.S. Patent Application No. 08/182,183, *supra*) described the truncation of mature GDNF at the carboxyl terminus by proteolytic processing of the Lys-Arg residues which are the sixth and fifth residues, respectively, from the carboxyl terminal end of mature GDNF (i.e., Lys¹²⁹-Arg¹³⁰ according to the amino acid residue numbering of Figure 1 (also as in SEQ ID NO:1 or SEQ ID NO:2). Such a truncation would eliminate two cysteine residues from the mature GDNF protein. This would be likely to result in improper folding of the protein, and therefore, would result in the formation of an inactive protein. In contrast, the X-[Cys⁴¹-Cys¹³³]-Y truncated GDNF protein products of the present invention retain the Cys¹³¹ and Cys¹³³ residues and are active proteins as determined by dopamine uptake assay.

In one embodiment of the present invention, preferred truncated GDNF protein products lack one or more deamidation sites. Such a lack of deamidation sites would result in the enhanced biochemical stability of the purified protein and a decrease in possible degradation products, thereby resulting in a more storage stable protein. An exemplary truncated GDNF protein product is the [Ser²⁶-Ile¹³⁴] truncated GDNF protein which lacks the sites which otherwise may lead to deamidation of the mature protein. Alternatively, the [Arg¹⁶-Ile¹³⁴] truncated GDNF protein would lack at least the first deamidation site otherwise present in the mature protein.

A currently preferred truncated GDNF protein product is [Arg³²-Ile¹³⁴] truncated GDNF protein. This truncated GDNF protein lacks the site at or near which proteolytic clipping of the mature protein occurs. Therefore, this truncated GDNF protein is expected to be resistant to the processing event which may also occur *in vivo*. Another currently preferred truncated GDNF protein product is the [Lys³⁷-Ile¹³⁴] truncated GDNF protein. This truncation would further reduce the pI of the truncated protein, as would other truncations in

which residues up to and including Gly⁴⁰ and Ile¹³⁴ are removed from the N- and C-terminals, respectively. The presently most preferred truncated GDNF protein products retain all of the cysteine residues found in mature GDNF protein, but lack any discernible sites for rapid proteolytic processing of the truncated GDNF protein during expression and manufacturing or following *in vivo* administration. These preferred proteins include the [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴], [Gln³⁴-Ile¹³⁴], [Arg³⁵-Ile¹³⁴], [Gly³⁶-Ile¹³⁴], [Lys³⁷-Ile¹³⁴], [Asn³⁸-Ile¹³⁴] and [Arg³⁹-Ile¹³⁴] truncated GDNF protein products.

Similar to the results previously described for mature GDNF by Lin et al. (U.S. Patent Application No. 07/855,413, *supra*), the truncated GDNF proteins of the present invention have demonstrated the ability to increase dopamine uptake by the embryonic precursors of the substantia nigra dopaminergic neurons. Bioassays of the truncated GDNF proteins are further described in Example 4, below.

The novel truncated GDNF proteins are typically isolated and purified to form truncated GDNF proteins which are substantially free from the presence of other (non-GDNF) proteinaceous materials. Preferably, the truncated GDNF protein products are about 80% free of other proteins which may be present due to the production technique used in the manufacture of the truncated GDNF protein product. More preferably, the truncated GDNF protein products are about 90% free of other proteins, particularly preferably, about 95% free of other proteins, and most preferably about >98% free of other proteins. In addition, the present invention furnishes the unique advantage of providing polynucleotide sequences for the manufacture of homogeneous truncated GDNF proteins. For example, the use of the polynucleotide sequence encoding the [Arg³²-Ile¹³⁴] truncated GDNF protein allows the recombinant production of the truncated GDNF protein in *E. coli* and other appropriate expression systems. In other words, the novel polynucleotides allow the production of truncated GDNF proteins which are not susceptible to proteolytic processing, or which have reduced susceptibility to such processing or other biochemical processing effects as described above. Thus, the novel polynucleotides make it easier to prepare and/or isolate single species truncated GDNF proteins, and therefore, the truncated GDNF proteins and/or products thereof do not contain or contain decreased amounts of the above-described mixture of hetero- and homodimers. It will be appreciated, however, that the final truncated GDNF protein products

may be combined with other factors, chemical compositions and/or suitable pharmaceutical formulation materials prior to administration, as described in further detail below.

In one aspect of the present invention, the truncated GDNF proteins are advantageously produced via recombinant techniques because they are capable of achieving comparatively higher amounts of protein at greater purity. Recombinant truncated GDNF protein forms include glycosylated and non-glycosylated forms of the protein, and protein expressed in bacterial, mammalian or insect cell systems. Alternatively, the truncated GDNF proteins may be chemically synthesized. Currently preferred production methods are described in greater detail below.

Truncated GDNF Variants and Derivatives

A. Truncated GDNF Variants

Another aspect of the present invention includes variants of truncated GDNF protein. The term "truncated GDNF protein products" as used herein includes variant proteins in which amino acids have been deleted from ("deletion variants"), inserted into ("addition variants"), or substituted for ("substitution variants"), residues within the amino acid sequence of naturally-occurring GDNF. Such variants are prepared by introducing appropriate nucleotide changes into the DNA encoding the protein or by *in vitro* chemical synthesis of the desired protein. It will be appreciated by those skilled in the art that many combinations of deletions, insertions, and substitutions can be made provided that the final protein possesses GDNF biological activity.

Mutagenesis techniques for the replacement, insertion or deletion of one or more selected amino acid residues are well known to one skilled in the art (e.g., U.S. Pat. No. 4,518,584, the disclosure of which is hereby incorporated by reference.) There are two principal variables in the construction of amino acid sequence variants: the location of the mutation site and the nature of the mutation. In designing truncated GDNF variants, the location of the mutation site and the nature of the mutation will depend on the biochemical characteristic(s) to be modified. The mutation sites can be modified individually or in series, e.g., by (1) substituting first with conservative amino acid choices and then with more radical selections depending upon the results achieved, (2) deleting the target amino acid residue, or (3) inserting amino acid residues adjacent to the located site.

Amino acid sequence deletions generally range from about 1 to 30 amino acid residues, more usually from about 1 to 10 residues, and typically from about 1 to 5 residues. For example, deletions in the "X" portion of the amino acid residues located N-terminally to Cys⁴¹ may range from approximately 1 to 30 residues, while deletions between the cysteine residues of [Cys⁴¹-Cys¹³³] are typically from about 1 to 5 residues, depending on the location, so as not to disrupt protein folding. Deletions within the truncated GDNF proteins may be made in regions of low homology with transforming growth factor-beta (TGF- β) family members. Deletions from truncated GDNF proteins in areas of substantial homology with other TGF- β family sequences will be more likely to modify the biological activity more significantly. The number of total deletions and/or consecutive deletions will be selected so as to preserve the tertiary structure of truncated GDNF protein in the affected domain, e.g., cysteine crosslinking.

Amino acid sequence additions may include amino- and/or carboxyl-terminal fusions ranging in length from one residue to one hundred or more residues, as well as internal intrasequence insertions of single or multiple amino acid residues. Internal additions may range generally from about 1 to 10 amino acid residues, more typically from about 1 to 5 amino acid residues, and usually from about 1 to 3 amino acid residues. As described above, the amino-terminus addition variants of the present invention are contemplated as including the addition of a methionine (for example, as an artifact of the direct expression of GDNF in bacterial recombinant cell culture) or a non-GDNF amino acid residue or sequence. Amino-terminus addition variants do not involve the addition of an amino acid residue(s) that would result in the reconstruction of the mature GDNF protein. A further example of a terminal insertion includes the fusion of a heterologous N-terminal signal sequence to the N-terminus to facilitate the secretion of protein from recombinant host cells. Such signal sequences generally will be obtained from, and thus be homologous to, the intended host cell species. Insertions or additions may also include amino acid sequences derived from the sequence of other neurotrophic factors.

Another group of variants are amino acid substitution variants. These variants have at least one amino acid residue in the truncated GDNF protein removed and a different residue inserted in its place. See, for example, Figure 5 wherein naturally occurring Asn²² was changed to Ser to facilitate further removal of the Met residue. Using the X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence representation and the present definition of truncated GDNF protein

products, such a truncated GDNF protein may be referred to either as a substitution variant Met-[Asn²²ΔSer²²-Ile¹³⁴] truncated GDNF protein or an addition variant Met-Ser-[Pro²³-Ile¹³⁴] truncated GDNF protein.

Substitution variants include allelic variants, which are characterized by

- 5 naturally-occurring nucleotide sequence changes in the species population that may or may not result in an amino acid change.

Specific mutations of the sequences of the truncated GDNF proteins may involve modifications of a glycosylation site (e.g., serine, threonine, or asparagine). The absence of glycosylation or only partial glycosylation may

- 10 result from amino acid substitution or deletion at any asparagine-linked glycosylation recognition site or at any site of the protein that is modified by the addition of an O-linked carbohydrate. An asparagine-linked glycosylation recognition site comprises a tripeptide sequence which is specifically recognized by appropriate cellular glycosylation enzymes. These tripeptide sequences are
- 15 either Asn-Xaa-Thr or Asn-Xaa-Ser, where Xaa can be any amino acid other than Pro. A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) result in non-glycosylation at the modified tripeptide sequence. Thus, the expression of appropriately altered nucleotide
- 20 sequences produces variants which are not glycosylated at that site. Alternatively, the sequence may be modified to add glycosylation sites to the truncated GDNF protein.

One method for identifying truncated GDNF amino acid residues or regions for mutagenesis is called "alanine scanning mutagenesis" as described by

- 25 Cunningham and Wells (*Science*, 244: 1081-1085, 1989). In this method, an amino acid residue or group of target residues are identified (e.g., charged residues such as Arg, Asp, His, Lys, and Glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to effect the interaction of the amino acids with the surrounding aqueous environment in
- 30 or outside the cell. Those domains demonstrating functional sensitivity to the substitutions then are refined by introducing additional or alternate residues at the sites of substitution. Thus, the site for introducing an amino acid sequence modification is predetermined, and to optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted and the
- 35 variants are screened for the optimal combination of desired activity and degree of activity.

The sites of greatest interest for substitutional mutagenesis include sites where the amino acids found in GDNF proteins from various species are substantially different in terms of side-chain bulk, charge, and/or hydrophobicity. Other sites of interest include those in which particular residues of GDNF-like proteins, obtained from various species, are identical. Such positions are generally important for the biological activity of a protein. Initially, these sites are modified by substitution in a relatively conservative manner. Such conservative substitutions are shown in Table 1 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes (exemplary substitutions) are introduced and/or other additions/deletions may be made, and the resulting products screened.

TABLE 1
Amino Acid Substitutions

<u>Original Residue</u>	<u>Preferred Substitutions</u>	<u>Exemplary Substitutions</u>
Ala (A)	Val	Val; Leu; Ile
Arg (R)	Lys	Lys; Gln; Asn
Asn (N)	Gln	Gln; His; Lys; Arg
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro	Pro
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; norleucine
Leu (L)	Ile	norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Leu	Leu; Val; Ile; Ala
Pro (P)	Gly	Gly
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; norleucine

Conservative modifications to the amino acid sequence (and the corresponding modifications to the encoding nucleic acid sequences) are expected

to produce truncated GDNF proteins having functional and chemical characteristics similar to those of the truncated GDNF proteins described in the Examples, below. In contrast, substantial modifications in the functional and/or chemical characteristics of truncated GDNF proteins may be accomplished by

5 selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the protein at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side chain

10 properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 4) basic: Asn, Gln, His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions may involve the exchange of a member of one of these classes for another. Such substituted residues may be introduced into regions of the truncated GDNF proteins that are homologous with other TGF- β proteins, or into the non-homologous regions of the protein.

20

B. Truncated GDNF Derivatives

Chemically modified derivatives of truncated GDNF or truncated GDNF variants may be prepared by one skilled in the art given the disclosures herein.

25 The chemical moieties most suitable for derivatization of truncated GDNF proteins include water soluble polymers. A water soluble polymer is desirable because the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Preferably, the polymer will be pharmaceutically acceptable for the preparation of a therapeutic product or composition. One skilled in the art will be able to select the desired polymer

30 based on such considerations as whether the polymer/protein conjugate will be used therapeutically, and if so, the desired dosage, circulation time, resistance to proteolysis, and other considerations. The effectiveness of the derivatization may be ascertained by administering the derivative, in the desired form (i.e., by

35 osmotic pump, or, more preferably, by injection or infusion, or further

formulated for oral, pulmonary or other delivery routes), and determining its effectiveness.

Suitable water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, monomethoxy-polyethylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, propylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), polyethylene glycol propionaldehyde, and mixtures thereof. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched.

The present invention particularly relates to truncated GDNF protein products involving truncated GDNF protein linked to at least one PEG molecule. In another aspect, the present invention relates to truncated GDNF protein attached to at least one PEG molecule via an acyl or alkyl linkage.

Pegylation may be carried out by any of the pegylation reactions known in the art. See, for example: *Focus on Growth Factors* 3(2): 4-10 (1992); EP 0 154 316; EP 0 401 384; and Malik et al., *Exp. Hematol.* 20: 1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive water soluble polymer. These preferred means for derivatization are discussed in greater detail, below. For the acylation reactions, the polymer(s) selected preferably have a single reactive ester group. For the reductive alkylation reactions, the polymer(s) selected preferably have a single reactive aldehyde group. In addition, the selected polymer may be modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled. Generally, the water soluble polymer will not be selected from naturally-occurring glycosyl residues since these are usually made more conveniently by mammalian recombinant expression systems.

Acylation

In the present invention, pegylation by acylation generally involves reacting an active ester derivative of polyethylene glycol with a truncated GDNF protein. Any known or subsequently discovered reactive PEG molecule may be used to carry out the pegylation process. A preferred activated PEG ester is PEG esterified to N-hydroxysuccinimide ("NHS"). As used herein, "acylation" is contemplated to include without limitation the following types of linkages between a truncated GDNF protein and a water soluble polymer such as PEG: amide, carbamate, urethane, and the like. See *Bioconjugate Chem.* 5: 133-140 (1994). Reaction conditions may be selected from any of those known in the pegylation art or those subsequently developed, but should avoid or limit exposure to reaction conditions such as temperatures, solvents, and pH levels that would inactivate the truncated GDNF protein to be modified.

Pegylation by acylation will generally result in a poly-pegylated truncated GDNF protein, wherein the lysine ϵ -amino groups are pegylated via an acyl linking group. Preferably, the connecting linkage will be an amide. Also preferably, the resulting product will be substantially only (e.g., $\geq 95\%$) mono-, di- or tri- pegylated. However, some conjugates with higher degrees of pegylation may be formed in amounts depending on the specific reaction conditions used. If desired, more purified pegylated conjugates may be prepared from the mixture by standard purification techniques, including, among others, dialysis, salting-out, ultrafiltration, ion-exchange chromatography, gel filtration chromatography and electrophoresis.

Alkylation

In the present invention, pegylation by alkylation generally involves reacting a terminal aldehyde derivative of PEG with a truncated GDNF protein in the presence of a reducing agent. Pegylation by alkylation can also result in a poly-pegylated truncated GDNF protein. In addition, one can manipulate the reaction conditions to favor pegylation substantially only at the α -amino group of the N-terminus of the protein (i.e., a mono-pegylated species). In either case of mono-pegylation or polypegylation, the PEG groups are preferably attached to the protein via a $-\text{CH}_2\text{-NH}-$ group. With particular reference to the $-\text{CH}_2-$ group, this type of linkage is referred to herein as an "alkyl" linkage.

Selective N-terminal chemical modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of

primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group-containing polymer is achieved. For example, one may selectively N-terminally pegylate the protein by performing the reaction at a pH which allows one to take advantage of the pK_a differences between the ϵ -amino group of the lysine residues and that of the α -amino group of the N-terminal residue of the protein. By such selective derivatization, attachment of a water soluble polymer to a protein is controlled: the conjugation with the polymer takes place predominantly at the N-terminus of the protein and no significant modification of other reactive groups, such as the lysine side chain amino groups, occurs. Using reductive alkylation, the water soluble polymer preferably has a single reactive aldehyde for coupling to the protein. Polyethylene glycol propionaldehyde, containing a single reactive aldehyde, may be used.

The present invention includes pegylated truncated GDNF proteins, wherein the PEG group(s) is (are) attached via acyl or alkyl groups. As discussed above, such truncated GDNF protein products may be mono-pegylated or poly-pegylated (e.g., containing 2-6, preferably 2-5, PEG groups). The PEG groups are generally attached to the protein at the α - or ϵ -amino groups of amino acids, but it is also contemplated that the PEG groups could be attached to any reactive group of to the protein, which is sufficiently reactive to become attached to a PEG group under suitable reaction conditions. Thus, polyethylene glycol may be covalently bound to a protein via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated PEG molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residue. Those having a free carboxyl group may include aspartic acid residues, glutamic acid residues, and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching PEG molecule(s). For therapeutic purposes, attachment at an amino group, such as attachment at the N-terminus or lysine group is typically preferred. Attachment at residues important for receptor binding should be avoided if receptor binding is desired.

In one aspect, the present invention provides for a substantially homogeneous preparation of mono-polymer/truncated GDNF protein conjugate

wherein a polymer molecule has been attached substantially only (i.e., $\geq 95\%$) in a single location. More specifically, if PEG is used, the present invention also provides for pegylated a truncated GDNF protein lacking possibly antigenic linking groups, and having the PEG molecule directly coupled to the truncated

5 GDNF protein.

In addition, derivatives may be prepared using glycosylated, non-glycosylated or de-glycosylated truncated GDNF proteins. Typically, non-glycosylated truncated GDNF proteins are used. For example, the prokaryote-expressed [Arg³²-Ile¹³⁴] truncated GDNF protein may be chemically

10 derivatized to include mono- or poly-, e.g., 2-4, PEG moieties, attached via an acyl or alkyl group):

In general, chemical derivatization may be performed under any suitable condition used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated truncated GDNF proteins will

15 generally comprise the steps of (a) reacting a truncated GDNF protein with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the truncated GDNF protein becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined case-

20 by-case based on known parameters and the desired result. For example, the larger the ratio of PEG:protein, the greater the percentage of poly-pegylated product. The optimum ratio (in terms of efficiency of reaction in that there is no excess unreacted protein or polymer) may be determined by factors such as the desired degree of derivatization (e.g., mono-, di-, tri-, etc.), the molecular

25 weight of the polymer selected, whether the polymer is branched or unbranched, and the reaction conditions used.

Reductive alkylation to produce a substantially homogeneous population of mono-polymer/truncated GDNF protein conjugate will generally comprise the steps of (a) reacting a truncated GDNF protein with a reactive PEG molecule

30 under reductive alkylation conditions, at a pH suitable to permit selective modification of the α -amino group at the amino terminus of the truncated GDNF protein, and (b) obtaining the reaction product(s).

For a substantially homogeneous population of mono-polymer/truncated GDNF protein conjugate, the reductive alkylation reaction conditions are those

35 which permit the selective attachment of the water soluble polymer moiety to the N-terminus of a truncated GDNF protein. Such reaction conditions generally

provide for pK_a differences between the lysine amino groups and the α -amino group at the N-terminus (the pK_a being the pH at which 50% of the amino groups are protonated and 50% are not). The pH also affects the ratio of polymer to protein to be used. In general, if the pH is lower, a larger excess of polymer

5 to protein will be desired (i.e., the less reactive the N-terminal α -amino group, the more polymer needed to achieve optimal conditions). If the pH is higher, the polymer:protein ratio need not be as large (i.e., more reactive groups are available, so fewer polymer molecules are needed). For purposes of the present invention, the pH will generally fall within the range of 3-9, preferably 3-6.

10 Another consideration is the molecular weight of the polymer. In general, the higher the molecular weight of the polymer, the fewer the number of polymer molecules which may be attached to the protein. Similarly, branching of the polymer should be taken into account when optimizing these parameters. Generally, the higher the molecular weight (or the more branches) the higher

15 the polymer:protein ratio. In general, for the pegylation reactions contemplated herein, the preferred average molecular weight is about 2 kDa to about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight). The preferred average molecular weight is about 5 kDa to about 50 kDa, particularly

20 preferably about 12 kDa to about 25 kDa. The ratio of water-soluble polymer to truncated GDNF protein will generally range from 1:1 to 100:1, preferably (for polypegylation) 1:1 to 20:1, and (for mono-pegylation) 1:1 to 5:1.

Using the conditions indicated above, reductive alkylation will provide for selective attachment of the polymer to any truncated GDNF protein having an

25 α -amino group at the amino terminus, and provide for a substantially homogenous preparation of mono-polymer/truncated GDNF protein conjugate. The term "mono-polymer/truncated GDNF protein conjugate" is used here to mean a derivative containing a single polymer molecule attached to a truncated GDNF protein. The mono-polymer/truncated GDNF protein conjugate preferably

30 will have a polymer molecule located at the N-terminus, but not on lysine amino side groups. The preparation will preferably be greater than 90% mono-polymer/truncated GDNF protein conjugate, and more preferably greater than 95% mono-polymer/truncated GDNF protein conjugate, with the remainder of observable proteins being unreacted (i.e., protein lacking the polymer moiety).

35 For reductive alkylation, the reducing agent should be stable in aqueous solution and preferably be able to reduce only the Schiff base formed in the

initial process of reductive alkylation. Exemplary reducing agents may be selected from the group consisting of sodium borohydride, sodium cyanoborohydride, dimethylamine borane, trimethylamine borane and pyridine borane. A particularly preferred reducing agent is sodium cyanoborohydride.

- 5 Other reaction parameters, such as solvent, reaction times, temperatures, etc., and means of purification of products, can be determined case-by-case based on commonly available information relating to derivatization of proteins with water soluble polymers.

- One may choose to prepare a mixture of polymer/protein conjugates by
10 acylation and/or alkylation methods, and the advantage provided herein is that one may select the proportion of mono-polymer/ protein conjugate to include in the mixture. Thus, if desired, one may prepare a mixture of protein having various numbers of polymer molecules attached thereto (i.e., di-, tri-, tetra-, etc.) and combine with the mono-polymer/protein conjugate material prepared
15 using the present methods, and have a mixture with a predetermined proportion of mono-polymer/protein conjugate.

Polynucleotides Encoding Truncated GDNF Proteins

- The present invention further provides novel polynucleotides which
20 encode truncated GDNF proteins. When used as a hybridization probe or amplification primer, the nucleic acid sequence will be substantially free from all other nucleic acid sequences. For use in recombinant protein expression, the nucleic acid sequence will generally be substantially free from nucleic acid sequences encoding other proteins, unless a fusion protein is intended. Based
25 upon the present description and using the universal codon table, one of ordinary skill in the art can readily determine all of the nucleic acid sequences which encode the amino acid sequences of truncated GDNF proteins. Presently preferred nucleic acid sequences include those polynucleotides encoding the [Arg¹⁶-Ile¹³⁴], [Ser²⁶-Ile¹³⁴], [Arg³²-Ile¹³⁴], and [Lys³⁷-Ile¹³⁴] truncated
30 GDNF proteins. Examples of a variety of polynucleotides are depicted in Figures 5, 6 and 7 as well as those portions of Figures 1, 3 and 4 which encode truncated GDNF proteins. It will also be appreciated by those skilled in the art that the novel polynucleotides which encode truncated GDNF proteins include those nucleic acid sequences encoding variant truncated GDNF proteins, whether man-
35 made or naturally occurring.

Recombinant expression techniques, conducted in accordance with the descriptions set forth below, may be followed to produce these polynucleotides and express the various truncated GDNF proteins. For example, by inserting a nucleic acid sequence which encodes a truncated GDNF protein into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate detection probes or amplification primers. Alternatively, a polynucleotide encoding a truncated GDNF protein can be inserted into an expression vector. By introducing the expression vector into an appropriate host, the desired truncated GDNF protein may be produced in large amounts.

As further described herein, there are numerous host/vector systems available for the propagation of nucleic acid sequences and/or the production of truncated GDNF proteins. These include, but are not limited to, plasmid, viral and insertional vectors, and prokaryotic and eukaryotic hosts. One skilled in the art can adapt a host/vector system which is capable of propagating or expressing heterologous DNA to produce or express the sequences of the present invention.

By means of such recombinant techniques, the truncated GDNF proteins of the present invention are readily produced in commercial quantities. Furthermore, it will be appreciated by those skilled in the art that, in view of the present disclosure, the novel nucleic acid sequences include degenerate nucleic acid sequences encoding the truncated GDNF proteins specifically set forth in the Figures, variants of such truncated GDNF proteins, and those nucleic acid sequences which hybridize, preferably under stringent hybridization conditions, to complements of these nucleic acid sequences (see, Maniatis et. al., *Molecular Cloning* (A Laboratory Manual); Cold Spring Harbor Laboratory, pages 387 to 389, 1982.) Exemplary stringent hybridization conditions are hybridization in 4 x SSC at 62-67°C, followed by washing in 0.1 x SSC at 62-67°C for approximately an hour. Alternatively, exemplary stringent hybridization conditions are hybridization in 45-55% formamide, 4 x SSC at 40-45°C. DNA sequences which hybridize to the complementary sequences for truncated GDNF protein under relaxed hybridization conditions and which encode a truncated GDNF protein of the present invention are also included herein. Examples of such relaxed stringency hybridization conditions are 4 x SSC at 45-55°C or hybridization with 30-40% formamide at 40-45°C.

Also provided by the present invention are recombinant DNA constructs involving vector DNA together with the DNA sequence encoding a truncated GDNF

protein. In such DNA constructs, the nucleic acid sequence encoding truncated GDNF protein (with or without signal peptides) is in operative association with a suitable expression control or regulatory sequence capable of directing the replication and/or expression of the truncated GDNF protein in a selected host.

5

Recombinant Expression of Truncated GDNF Protein

Preparation of Polynucleotides encoding Truncated GDNF

A nucleic acid sequence encoding truncated GDNF, or a mature GDNF starting material, can readily be obtained in a variety of ways, including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening, and/or PCR amplification of cDNA. These methods and others useful for isolating such nucleic acid sequences are set forth, for example, by Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), by Ausubel *et al.*, eds (*Current Protocols in Molecular Biology*, Current Protocols Press, 1994), and by Berger and Kimmel (*Methods in Enzymology: Guide to Molecular Cloning Techniques*, vol. 152, Academic Press, Inc., San Diego, CA, 1987). Preferred nucleic acid sequences encoding GDNF are mammalian sequences.

Chemical synthesis of a nucleic acid sequence which encodes a truncated GDNF protein can also be accomplished using methods well known in the art, such as those set forth by Engels *et al.* (*Angew. Chem. Intl. Ed.*, 28:716-734, 1989). These methods include, *inter alia*, the phosphotriester, phosphoramidite and H-phosphonate methods of nucleic acid sequence synthesis. The nucleic acid sequence encoding the truncated GDNF protein will be several hundred base pairs (bp) or nucleotides in length. Large nucleic acid sequences, for example those larger than about 100 nucleotides in length, can be synthesized as several fragments. The fragments can then be ligated together to form a nucleic acid sequence encoding truncated GDNF protein. A preferred method is polymer-supported synthesis using standard phosphoramidite chemistry.

Alternatively, a suitable nucleic acid sequence may be obtained by screening an appropriate cDNA library (i.e., a library prepared from one or more tissue source(s) believed to express the protein) or a genomic library (a library prepared from total genomic DNA). The source of the cDNA library is typically a tissue from any species that is believed to express GDNF in reasonable quantities. The source of the genomic library may be any tissue or tissues from any mammalian or other species believed to harbor a gene encoding

GDNF or a GDNF homologue. The library can be screened for the presence of the GDNF cDNA/gene using one or more nucleic acid probes (oligonucleotides, cDNA or genomic DNA fragments that possess an acceptable level of homology to the GDNF or GDNF homologue cDNA or gene to be cloned) that will hybridize

5 selectively with GDNF or GDNF homologue cDNA(s) or gene(s) present in the library. The probes typically used for such library screening usually encode a small region of the GDNF DNA sequence from the same or a similar species as the species from which the library was prepared. Alternatively, the probes may be degenerate, as discussed herein.

10 Library screening is typically accomplished by annealing the oligonucleotide probe or cDNA to the clones in the library under conditions of stringency that prevent non-specific binding but permit binding of those clones that have a significant level of homology with the probe or primer. Typical hybridization and washing stringency conditions depend in part on the size (i.e.,
15 number of nucleotides in length) of the cDNA or oligonucleotide probe, and whether the probe is degenerate. The probability of obtaining a clone(s) is also considered in designing the hybridization solution (i.e., whether a cDNA or genomic library is being screened; if it is a cDNA library, the probability that the cDNA of interest is present at a high level).

20 Where DNA fragments (such as cDNAs) are used as probes, typical hybridization conditions include those as set forth in Ausubel et al., *eds., supra*. After hybridization, the blot containing the library is washed at a suitable stringency, depending on several factors such as probe size, expected homology of probe to clone, type of library being screened, number of clones being screened,
25 and the like. Examples of stringent washing solutions (which are usually low in ionic strength and are used at relatively high temperatures) are as follows. One such stringent wash is 0.015 M NaCl, 0.005 M NaCitrate and 0.1% SDS at 55-65°C. Another such stringent buffer is 1 mM Na₂EDTA, 40 mM NaHPO₄, pH 7.2, and 1% SDS at about 40-50°C. One other stringent wash is 0.2 X SSC and
30 0.1% SDS at about 50-65°C.

There are also exemplary protocols for stringent washing conditions where oligonucleotide probes are used to screen cDNA or genomic libraries. For example, a first protocol uses 6 X SSC with 0.05 percent sodium pyrophosphate at a temperature of between about 35 and 62°C, depending on the length of the
35 probe. For example, 14 base probes are washed at 35-40°C, 17 base probes at 45-50°C, 20 base probes at 52-57°C, and 23 base probes at 57-63°C. The

temperature can be increased 2-3°C where the background non-specific binding appears high. A second protocol uses tetramethylammonium chloride (TMAC) for washing. One such stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2% SDS.

5 Another suitable method for obtaining a nucleic acid sequence encoding a GDNF protein is the polymerase chain reaction (PCR). In this method, poly(A)+RNA or total RNA is extracted from a tissue that expresses GDNF. cDNA is then prepared from the RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of the GDNF cDNA
10 (oligonucleotides), are then added to the cDNA along with a polymerase such as *Taq* polymerase, and the polymerase amplifies the cDNA region between the two primers.

Where the method of choice for preparing the nucleic acid sequence encoding the desired truncated GDNF protein requires the use of oligonucleotide
15 primers or probes (e.g., PCR, cDNA or genomic library screening), the oligonucleotide sequences selected as probes or primers should be of adequate length and sufficiently unambiguous so as to minimize the amount of non-specific binding that will occur during library screening or PCR amplification. The actual sequence of the probes or primers is usually based on conserved or highly
20 homologous sequences or regions from the same or a similar gene from another organism. Optionally, the probes or primers can be fully or partially degenerate, i.e., contain a mixture of probes/primers, all encoding the same amino acid sequence, but using different codons to do so. An alternative to preparing degenerate probes is to place an inosine in some or all of those codon
25 positions that vary by species. The oligonucleotide probes or primers may be prepared by chemical synthesis methods for DNA as described above.

Truncated GDNF proteins based on these nucleic acid sequences encoding GDNF, as well as mutant or variant sequences thereof, are also contemplated as within the scope of the present invention. As described above, a mutant or
30 variant sequence is a sequence that contains one or more nucleotide substitutions, deletions, and/or insertions as compared to the wild type sequence and that results in the expression of amino acid sequence variations as compared to the wild type amino acid sequence. In some cases, naturally occurring GDNF amino acid mutants or variants may exist, due to the existence of natural allelic
35 variation. Truncated GDNF proteins based on such naturally occurring mutants or variants are also within the scope of the present invention. Preparation of

synthetic mutant sequences is also well known in the art, and is described for example in Wells et al. (*Gene*, 34:315, 1985) and in Sambrook et al., *supra*.

Vectors

- 5 The cDNA or genomic DNA encoding a truncated GDNF protein is inserted into a vector for further cloning (amplification of the DNA) or for expression. Suitable vectors are commercially available, or the vector may be specially constructed. The selection or construction of the appropriate vector will depend on 1) whether it is to be used for DNA amplification or for DNA expression, 2) 10 the size of the DNA to be inserted into the vector, and 3) the host cell (e.g., mammalian, insect, yeast, fungal, plant or bacterial cells) to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and its compatibility with the intended host cell. The vector components generally include, but are not 15 limited to, one or more of the following: a signal sequence, an origin of replication, one or more selection or marker genes, enhancer elements, promoters, a transcription termination sequence, and the like. These components may be obtained from natural sources or synthesized by known procedures. The vectors of the present invention involve a nucleic acid sequence 20 which encodes the truncated GDNF protein of interest operatively linked to one or more of the following expression control or regulatory sequences capable of directing, controlling or otherwise effecting the expression of the truncated GDNF protein by a selected host cell.

25 Signal Sequence

- The signal sequence may be a component of the vector, or it may be a part of GDNF DNA that is inserted into the vector. The native GDNF DNA encodes a signal sequence at the amino terminus of the protein that is cleaved during post-translational processing of the protein to form the mature GDNF protein. 30 Included within the scope of this invention are truncated GDNF polynucleotides with the native signal sequence and other pre-pro sequences as well as truncated GDNF polynucleotides wherein the native signal sequence is deleted and replaced with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, 35 by the host cell. For prokaryotic host cells that do not recognize and process the native GDNF signal sequence, the signal sequence is substituted by a prokaryotic

signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native GDNF signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

Origin of Replication

Expression and cloning vectors generally include a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. In cloning vectors, this sequence is typically one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeasts, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria and various origins (e.g., SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

Selection Gene

The expression and cloning vectors typically contain a selection gene. This gene encodes a "marker" protein necessary for the survival or growth of the transformed host cells when grown in a selective culture medium. Host cells that were not transformed with the vector will not contain the selection gene, and therefore, they will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline; (b) complement auxotrophic deficiencies; or (c) supply critical nutrients not available from the culture medium.

Other selection genes may be used to amplify the gene which will be expressed. Amplification is the process wherein genes which are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells.

Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell

transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of the marker present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes truncated GDNF. As a result, increased quantities of truncated GDNF are synthesized from the amplified DNA.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate, a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is used is the Chinese hamster ovary cell line deficient in DHFR activity (see, for example, Urlaub and Chasin, *Proc. Natl. Acad. Sci., USA* 77(7): 4216-4220 (1980)). The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA present in the expression vector, such as the DNA encoding a truncated GDNF protein.

Promoter

The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to the nucleic acid sequence encoding the truncated GDNF protein. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular nucleic acid sequence, such as that encoding truncated GDNF. Promoters are conventionally grouped into one of two classes, inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. A large number of promoters, recognized by a variety of potential host cells, are well known. These promoters are operably linked to the DNA encoding truncated GDNF by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native GDNF promoter sequence may be used to direct amplification and/or expression of truncated GDNF DNA. A heterologous promoter is preferred, however, if it permits greater transcription and higher

yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase, a tryptophan (trp) promoter system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their nucleotide sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence(s), using linkers or adaptors as needed to supply any required restriction sites.

Suitable promoting sequences for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, e.g., heat-shock promoters and the actin promoter. A currently used promoter in the production of GDNF proteins in CHO cells is SR α . See Takebe et al., *Mol. Cell. Biol.* 8(1): 466-472 (1988). A suitable expression vector is pDSR α 2, which is further described below.

Enhancer Element

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA sequence encoding a truncated GDNF protein of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about from 10-300 bp in length, that act on the promoter to increase its transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to truncated GDNF DNA, it is typically located at a site 5' from the promoter.

Transcription Termination

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and occasionally 3' untranslated regions of eukaryotic DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding truncated GDNF.

The construction of suitable vectors containing one or more of the above-listed components together with the desired truncated GDNF coding sequence is accomplished by standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the desired order to generate the plasmids required. To confirm that the correct sequences have been constructed, the ligation mixtures may be used to transform *E. coli*, and successful transformants may be selected by known techniques, such as ampicillin or tetracycline resistance as described above. Plasmids from the transformants are then prepared, analyzed by restriction endonuclease digestion, and/or sequenced to confirm the presence of the desired construct.

Vectors that provide for the transient expression of DNA encoding truncated GDNF in mammalian cells may also be used. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of the desired protein encoded by the expression vector. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of proteins encoded by cloned DNAs, as well as for the rapid screening of such proteins for desired biological or physiological properties. Thus, transient expression systems are particularly useful in identifying variants of the protein.

Selection and Transformation of Host Cells

Host cells (e.g., bacterial, mammalian, insect, yeast, or plant cells) transformed with nucleic acid sequences for use in expressing a recombinant truncated GDNF protein are also provided by the present invention. The transformed host cell is cultured under appropriate conditions permitting the

expression of the nucleic acid sequence. The selection of suitable host cells and methods for transformation, culture, amplification, screening and product production and purification are well known in the art. See for example, Gething and Sambrook, *Nature* 293: 620-625 (1981), or alternatively, Kaufman et al.,
5 *Mol. Cell. Biol.*, 5 (7): 1750-1759 (1985) or Howley et al., U.S. Pat. No. 4,419,446. Truncated GDNF may be expressed in *E. coli* in accordance with the description of Lin et al. (U.S. Patent Application No. 07/855,413; Application No. PCT/US92/07888; WO 93/06116) which involved the expression of mature GDNF. Other exemplary materials and methods are discussed in further
10 detail below. The transformed host cell is cultured in a suitable medium, and the expressed factor is then optionally recovered, isolated and purified from the culture medium (or from the cell, if expressed intracellularly) by an appropriate means known to those skilled in the art.

Suitable host cells for cloning or expressing the vectors herein are the
15 prokaryote, yeast, or higher eukaryote cells as described above. Prokaryotic host cells include, but are not limited to, eubacteria, such as Gram-negative or Gram-positive organisms, for example, *E. coli*, *Bacilli* such as *B. subtilis*, *Pseudomonas* species such as *P. aeruginosa*, *Salmonella typhimurium*, or *Serratia marcescans*. Alternatively, *in vitro* methods of cloning, e.g., PCR or
20 other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotic host cells, eukaryotic microbes such as filamentous fungi or yeast may be suitable hosts for the expression of truncated GDNF proteins. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms, but a number of
25 other genera, species, and strains are well known and commonly available.

Suitable host cells for the expression of glycosylated truncated GDNF protein are derived from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture might be used, whether such culture involves vertebrate or invertebrate cells, including plant and insect cells. Vertebrate cells are
30 generally used as the propagation of vertebrate cells in culture (tissue culture) is a well known procedure. Examples of useful mammalian host cell lines include, but are not limited to, monkey kidney CV1 line transformed by SV40 (COS-7), human embryonic kidney line (293 or 293 cells subcloned for growth
35 in suspension culture), baby hamster kidney cells, and Chinese hamster ovary cells. Other suitable mammalian cell lines include but are not limited to, HeLa,

mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *Streptomyces spp.* and the like may also be employed.

Presently preferred host cells for producing truncated GDNF proteins are bacterial cells (e.g., *Escherichia coli*) and mammalian cells (such as Chinese hamster ovary cells, COS cells, etc.)

The host cells are transfected and preferably transformed with the above-described expression or cloning vectors and cultured in a conventional nutrient medium. The medium may be modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Transfection and transformation are performed using standard techniques which are well known to those skilled in the art and which are selected as appropriate to the host cells involved. For example, for mammalian cells without cell walls, the calcium phosphate precipitation method may be used. Electroporation, micro injection and other known techniques may also be used.

Culturing the Host Cells

Transformed cells used to produce truncated GDNF proteins of the present invention are cultured in suitable media. The media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as gentamicin), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or other energy source. Other supplements may also be included, at appropriate concentrations, as will be appreciated by those skilled in the art. Suitable culture conditions, such as temperature, pH, and the like, are also well known to those skilled in the art for use with the selected host cells.

It is also possible that truncated GDNF proteins may be produced by homologous recombination, or with recombinant production methods utilizing control elements introduced into cells already containing DNA encoding GDNF. Homologous recombination is a technique originally developed for targeting genes

- to induce or correct mutations in transcriptionally active genes (Kucherlapati, *Prog. in Nucl. Acid Res. and Mol. Biol.* 36:301 (1989)). The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas et al., *Cell.* 44:419-428, 1986;
- 5 Thomas and Capecchi, *Cell.* 51:503-512, 1987; Doetschman et al., *Proc. Natl. Acad. Sci.* 85:8583-8587, 1988) or to correct specific mutations within defective genes (Doetschman et al., *Nature.* 330:576-578, 1987). Exemplary homologous recombination techniques are described in U.S. 5,272,071 (EP 91 90 3051, EP Publication No. 505 500; PCT/US90/07642, International
- 10 Publication No. WO 91/09955) the disclosure of which is hereby incorporated by reference.

- Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is DNA that is complementary
- 15 (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize and therefore recombine with other pieces of endogenous DNA through shared homologous regions. If this
- 20 complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence of DNA, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the new sequence of DNA to serve as the template. Thus, the transferred DNA is incorporated into the genome.

- 25 If the sequence of a particular gene is known, such as the nucleic acid sequence of GDNF, the pre-pro sequence or expression control sequence, a piece of DNA that is complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a
- 30 targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during DNA replication, this piece of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be backstitched into the newly synthesized daughter strand of DNA.

- 35 In the present invention, attached to these pieces of targeting DNA are regions of DNA which may interact with the expression of a GDNF protein. For

example, a promoter/enhancer element, a suppresser, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired truncated GDNF. The control element does not encode

- 5 truncated GDNF, but instead controls a portion of the DNA present in the host cell genome. Thus, the expression of truncated GDNF proteins may be achieved not by transfection of DNA that encodes the truncated GDNF gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene
10 sequence with recognizable signals for transcription of a truncated GDNF protein.

In accordance with the present invention, homologous recombination methods may also be used to modify a cell that contains a normally transcriptionally silent GDNF gene to produce a cell which expresses GDNF. The GDNF protein may then be processed to form a truncated GDNF protein(s).

15

Truncated GDNF Pharmaceutical Compositions

- Truncated GDNF protein product pharmaceutical compositions typically include a therapeutically effective amount of a truncated GDNF protein product in admixture with one or more pharmaceutically and physiologically acceptable
20 formulation materials. Suitable formulation materials include, but are not limited to, antioxidants, preservatives, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers, bulking agents, buffers, delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. For
25 example, a suitable vehicle may be water for injection, physiological saline solution, or artificial cerebrospinal fluid (CSF), possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles.

- The primary solvent in a vehicle may be either aqueous or non-aqueous
30 in nature. In addition, the vehicle may contain other pharmaceutically-acceptable excipients for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the vehicle may contain still other pharmaceutically-acceptable excipients for modifying or maintaining the stability, rate of
35 dissolution, or rate of release of truncated GDNF protein product, or for promoting the absorption or penetration of truncated GDNF protein product

across the blood-brain barrier. Such excipients are those substances usually and customarily employed to formulate dosages for parenteral administration in either unit dose or multi-dose form or for direct infusion into the CSF by continuous or periodic infusion from an implanted pump.

5 Once the therapeutic composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready to use form or in a form, e.g., lyophilized, requiring reconstitution prior to administration.

10 The optimal pharmaceutical formulation will be determined by one skilled in the art depending upon the route of administration and desired dosage. See for example, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton, PA 18042) pages 1435-1712 the disclosure of which is hereby incorporated by reference. The composition may also involve particulate
15 preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hylauronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the present proteins and derivatives.

20 Other effective administration forms, such as parenteral slow-release formulations, inhalant mists, orally active formulations, or suppositories, are also envisioned. Current truncated GDNF protein product pharmaceutical compositions are formulated for parenteral administration, e.g., intracerebroventricular injection. Such parenterally administered therapeutic
25 compositions are typically in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising truncated GDNF protein product in a pharmaceutically acceptable vehicle. One preferred vehicle is physiological saline.

 It is also contemplated that certain formulations containing truncated
30 GDNF protein product are to be administered orally. Truncated GDNF protein product which is administered in this fashion may be encapsulated and may be formulated with or without those carriers customarily used in the compounding of solid dosage forms. The capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is
35 maximized and pre-systemic degradation is minimized. Additional excipients may be included to facilitate absorption of truncated GDNF protein product.

Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

5 Administration of Truncated GDNF Protein Product

The truncated GDNF protein product may be administered parenterally via a subcutaneous, intramuscular, intravenous, transpulmonary, transdermal, intrathecal or intracerebral route. Protein growth factors that do not cross the blood-brain barrier may be given directly intracerebrally or otherwise in association with other elements that will transport them across the barrier. It is preferred that the truncated GDNF protein product is administered intracerebroventricularly or into the brain or spinal cord subarachnoid space. Truncated GDNF protein product may also be administered intracerebrally directly into the brain parenchyma. Slow-releasing implants in the brain containing the neurotrophic factor embedded in a biodegradable polymer matrix can also deliver truncated GDNF protein product. Truncated GDNF protein product may be administered extracerebrally in a form that has been modified chemically or packaged so that it passes the blood-brain barrier, or it may be administered in connection with one or more agents capable of promoting penetration of truncated GDNF protein product across the barrier. For example, a conjugate of NGF and monoclonal anti-transferrin receptor antibodies has been shown to be transported to the brain via binding to transferrin receptors. To achieve the desired dose of truncated GDNF protein product, repeated daily or less frequent injections may be administered, or truncated GDNF protein product may be infused continuously or periodically from a constant- or programmable-flow implanted pump. The frequency of dosing will depend on the pharmacokinetic parameters of the truncated GDNF protein product as formulated, and the route of administration.

Regardless of the manner of administration, the specific dose is typically calculated according to body weight or body surface area. For diseases involving the brain, the specific dose is typically calculated according to the approximate brain weight of the patient, which also may be estimated based on body weight or body surface area. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art, especially in light of the dosage information and assays disclosed herein.

Appropriate dosages may be ascertained through use of the established assays for determining dosages utilized in conjunction with appropriate dose-response data. The final dosage regimen involved in a method of treating a specific condition will be determined by the attending physician, considering various factors which

5 modify the action of drugs, e.g., the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors.

Truncated GDNF protein product of the present invention may also be employed, alone or in combination with other growth factors in the treatment of

10 nerve disease. For example, truncated GDNF protein product may be used in treating some forms of nerve disease in combination with nerve growth factor. In addition, other factors or other molecules, including chemical compositions, may be employed together with truncated GDNF protein product. In the treatment of Parkinson's Disease, it is contemplated that truncated GDNF protein product

15 be used by itself or in conjunction with the administration of Levodopa, wherein the truncated GDNF would enhance the production of endogenous dopamine and the neuronal uptake of the increased concentration of dopamine.

As stated above, it is also contemplated that additional neurotrophic or neuron nurturing factors will be useful or necessary to treat some neuronal cell

20 populations or some types of injury or disease. Other factors that may be used in conjunction with truncated GDNF include, but are not limited to: mitogens such as insulin, insulin-like growth factors, epidermal growth factor, vasoactive growth factor, pituitary adenylate cyclase activating polypeptide, interferon and somatostatin; neurotrophic factors such as brain derived neurotrophic factor,

25 neurotrophin-3, neurotrophin-4/5, neurotrophin-6, insulin-like growth factor, ciliary neurotrophic factor, acidic and basic fibroblast growth factors, fibroblast growth factor-5, transforming growth factor- β , cocaine-amphetamine regulated transcript (CART) and mature GDNF; and other growth factors such as epidermal growth factor, leukemia inhibitory factor,

30 interleukins, interferons, and colony stimulating factors; as well as molecules and materials which are the functional equivalents to these factors.

It is envisioned that the continuous administration or sustained delivery of a truncated GDNF protein product may be advantageous for a given treatment. While continuous administration may be accomplished via a mechanical means,

35 such as with an infusion pump, it is contemplated that other modes of continuous or near continuous administration may be practiced. For example, chemical

derivatization may result in sustained release forms of the protein which have the effect of continuous presence in the blood stream, in predictable amounts, based on a determined dosage regimen. Thus, truncated GDNF protein products include truncated GDNF protein derivatized to effectuate such continuous administration.

Truncated GDNF protein cell therapy, e.g., intracerebral implantation of cells producing truncated GDNF protein, is also contemplated. This embodiment of the present invention may include implanting into patients cells which are capable of synthesizing and secreting a biologically active form of truncated GDNF protein. Such truncated GDNF protein producing-cells may be cells which do not normally produce a neurotrophic factor but have been modified to produce truncated GDNF, or they could be cells whose ability to produce GDNF has been augmented by transformation with a polynucleotide suitable for the expression and secretion of truncated GDNF protein. In order to minimize a potential immunological reaction in patients from administering GDNF of a foreign species, it is preferred that the cells be of human origin and produce truncated human GDNF protein.

Implanted cells may be encapsulated to avoid infiltration of the cells into brain tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semi-permeable polymeric enclosures or membranes to allow release of a truncated GDNF protein product, but that prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue. Alternatively, the patient's own cells, transformed *ex vivo* to produce truncated GDNF, could be implanted directly into the patient without such encapsulation.

The methodology for the membrane encapsulation of living cells is familiar to those of ordinary skill in the art, and the preparation of the encapsulated cells and their implantation in patients may be accomplished. See for example, U.S. Patent Numbers 4,892,538; 5,011,472; and 5,106,627, the disclosures of which are hereby incorporated by reference. A system for encapsulating living cells is also described in PCT Application WO 91/10425 of Aebischer et al., specifically incorporated herein by reference. See also, PCT Application WO 91/10470 of Aebischer et al.; Winn et al., *Exper. Neurol.*, 113:322-329, 1991; Aebischer et al., *Exper. Neurol.*, 111:269-275, 1991; Tresco et al., *ASAIO*, 38:17-23, 1992, the disclosures of which are hereby incorporated by reference.

Truncated GDNF protein gene therapy *in vivo* is also envisioned, wherein a nucleic acid sequence encoding a truncated GDNF protein is introduced directly into the patient. For example, a nucleic acid sequence encoding a truncated GDNF protein is introduced into target cells via local injection of a nucleic acid

5 construct with or without an appropriate delivery vector, such as an adeno-associated virus vector. Alternative viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus and papilloma virus vectors. Physical transfer may be achieved *in vivo* by local injection of the desired nucleic acid construct or other appropriate delivery vector containing the

10 desired nucleic acid sequence, liposome-mediated transfer, direct injection (naked DNA), receptor-mediated transfer (ligand-DNA complex), or microparticle bombardment (gene gun).

It should be noted that the truncated GDNF protein product formulations described herein may be used for veterinary as well as human applications and

15 that the term "patient" should not be construed in a limiting manner. In the case of veterinary applications, the dosage ranges should be the same as specified above.

As a means of further characterizing truncated GDNF proteins of the present invention, antibodies can be developed which bind to the truncated GDNF

20 protein such as to epitopes within the X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence. One of ordinary skill in the art can use well-known, published procedures to obtain monoclonal and polyclonal antibodies, or recombinant antibodies, which specifically recognize and bind to the various proteins encoded by the amino acid sequences of the present invention. Such antibodies may then

25 be used to purify and characterize truncated GDNF protein. Alternatively, the antibodies may be used as therapeutical inhibitors of the proteins to which they are directed.

Other aspects and advantages of the present invention will be understood

30 upon consideration of the following illustrative examples. Example 1 addresses the expression of mature GDNF in a mammalian cell system and the preparation of truncated GDNF protein. Example 2 addresses the expression of mature GDNF in a bacterial cell system. Example 3 addresses the expression of various truncated GDNF proteins in a bacterial cell system. Example 4 compares the

35 biological activity of the mature GDNF protein and truncated GDNF protein in an assay for dopaminergic neuron neurotrophic activity.

EXAMPLES

5

Example 1

Expression of Mature Human GDNF in CHO Cells
and the Purification of CHO cell-derived Truncated GDNF Protein

Materials

10

The following materials are used in the expression of human GDNF in dihydrofolate reductase-deficient CHO cells (CHO⁻ cells; for example, as described by Urlaub and Chasin, *Proc. Natl. Acad. Sci., USA* 77(7): 4216-4220 (1980)).

15

CHO⁻ medium contained: Dulbecco's Modified Eagle's Medium (DMEM) - high glucose (Gibco/BRL); 5% fetal bovine serum (HyClone); MEM non-essential amino acids (1%) (Gibco/BRL); hypoxanthine/thymidine (1%) (Gibco/BRL); and glutamine/penicillin/streptomycin (1%) (Irvine Scientific).

20

Selective medium contained: DMEM (high glucose); 5% dialyzed fetal bovine serum (HyClone); MEM non-essential amino acids; and glutamine/penicillin/streptomycin.

2X HEPES-buffered saline (HBS) contained: 280 mM NaCl; 10 mM KCl; 1.5 mM Na₂HPO₄; 12 mM dextrose; and 50 mM HEPES.

Tris-buffered saline plus Tween (TBST) contained: 137 mM NaCl; 20 mM Tris/HCl pH 7.5; and 0.1% Tween-20.

25

Methods

Transfection and Selection

CHO⁻ cells (passage 20) were seeded into 60mm tissue culture dishes (Falcon) at a density of 8×10^5 cells per dish in CHO⁻ growth medium. On the following day, about three hours prior to transfection, the medium on the cells was replaced with fresh medium.

35

Plasmid constructs containing the appropriate GDNF cDNA were prepared using well known techniques. For example, the plasmid construct pDSRα2 was prepared substantially in accordance with the process described in the co-owned and copending U. S. Patent Application Serial Number 501,904 filed March 29, 1990 (also see, European Patent Application No. 90305433, Publication No.

EP 398 753, filed May 18, 1990 and WO 90/14363 (1990), the disclosures of which are hereby incorporated by reference. An exemplary plasmid map which illustrates the structural organization of the vector is depicted in Figure 2. It will be appreciated by those skilled in the art that a variety of nucleic acid sequences encoding the mature GDNF protein, such as the sequences depicted in Figures 1, 3 and 4, may also be used.

A HindIII- XbaI DNA fragment containing the human GDNF coding sequences and the consensus Kozak sequences, CCACC(ATG) was retrieved by restriction enzyme digestion from a pcDNA3 based expression vector (Invitrogen, San Diego, CA). The DNA fragment was directly cloned into HindIII/XbaI cut pDSR α 2. The resulting plasmid was called pSW5. The plasmid DNA of pSW5 was linearized at the PvuI site prior to transfection.

pDSR α 2 (Figure 2) is a derivative of the plasmid pCD (Okayama & Berg, *Mol. Cell Biol.* 3: 280-289, 1983) with three main modifications: (i) the SV40 polyadenylation signal has been replaced with the signal from the α -subunit of bovine follicular stimulating hormone, α -bFSH (Goodwin et al., *Nucleic Acids Res.* 11: 6873-6882, 1983); (ii) a mouse dihydrofolate reductase minigene (Gasser et al., *Proc. Natl. Acad. Sci.* 79: 6522-6526, 1982) has been inserted downstream from the expression cassette to allow selection and amplification of the transformants; and (iii) a 267 bp fragment containing the "R-element" and part of the "U5" sequences of the long terminal repeat (LTR) of human T-cell leukemia virus type I (HTLV-I) has been cloned and inserted between the SV40 promoter and the splice signals as described previously (Takebe et al., *Mol. Cell Biol.* 8: 466-472, 1988).

Solutions of DNA were prepared containing a final concentration of 3.0 μ g/dish GDNF-plasmid DNA, 7.0 μ g/dish mouse kidney genomic carrier DNA (Clontech), 25 μ l/dish 2.5M CaCl₂, and sterile distilled water to a final volume of 250 μ l/dish. DNA solutions containing pDSR α 2 vector DNA or carrier DNA alone were similarly prepared as positive and negative controls, respectively. The DNA solutions were added dropwise to an equal volume of 2X HEPES-buffered saline while passing air bubbles through the solution. The DNA/HBS solutions were incubated at room temperature for 30 minutes.

The medium was removed from the CHO^d- cell cultures, and 500 μ l of the DNA solutions were added per dish. The dishes were incubated at room temperature for 30 minutes, after which time CHO^d- medium (5.0 ml.) was added to each dish. The dishes were then incubated at 37°C overnight.

On the following day the medium was replaced with fresh CHO^d medium. The next day, when the cells had reached confluence, the cultures were trypsinized and replated in 100 mm dishes (Falcon) at a ratio of 1x60 mm dish to 8x100 mm dishes. Cells were replated in selective medium. The cultures were re-fed with fresh medium every two to three days.

After 15 days, colonies of transfected cells were isolated using glass cloning cylinders, trypsinized, and replated into 24-well dishes (Falcon). A total of 40 colonies was isolated from the GDNF/pSW5-transfected cells. The remaining cells on the dishes were trypsinized, pooled, and replated into two 100 mm dishes (one pool for each DNA construct).

Screening of transfected cells:

The 24-well and pool cultures were grown to confluence, at which time the growth medium was removed and replaced with serum-free medium (400 µl/well or 4 ml/dish). Cells were incubated for 48 hours, and the conditioned medium was harvested. The conditioned medium samples were analyzed for GDNF protein expression by western blot. Aliquots of conditioned medium (20 µl or 40 µl) were diluted with electrophoresis sample buffer (with or without β-mercaptoethanol). Samples containing β-mercaptoethanol were boiled for three minutes (reducing conditions). Both reduced and non-reduced samples were run on 16% Tris-glycine gels (Novex). Gels were electroblotted onto nitrocellulose filters (Schleicher and Schuell BA-83, 0.2µ). The blots were rinsed with TBST and then incubated in a blocking solution of 5% dried milk (Carnation) in TBST for 30 minutes at room temperature. Blots were then treated with GDNF antiserum (rabbit polyclonal antisera raised against *E. coli*-derived GDNF; 1:1000 in 5% milk/TBST) for one hour at room temperature. The blots were then rinsed with TBST and washed 1x10 minutes and 2x5 minutes with 1% milk/TBST. They were then treated with anti-rabbit Ig-horse radish peroxidase-conjugated secondary antibody (1:15,000 in 1% milk/TBST) for 20 minutes. Blots were rinsed and washed with TBST 1x20 minutes and 2x10 minutes, followed by treatment with ECL reagents (Amersham) for one minute and exposure to Hyperfilm-ECL (Amersham).

The following process describes the purification of CHO-expressed GDNF and a CHO-derived clipped GDNF homodimer, from one liter of conditioned media. Because of significant protease action in the CHO medium, clipping the chain at

residue 31, the procedure may include the use of a protease inhibitor during purification.

Step 1. Bead Chromatography:

5 Serum free conditioned media was made 20 mM 2-[N-Morpholino]
ethane sulfonate (MES), pH 6.0, by adding one fiftieth volume of 1 M MES, pH
6.0. Twenty five milliliters of SP Sepharose Big Bead resin (Pharmacia),
equilibrated with 20 mM MES, pH 6.0, was added and stirred at 4°C for one hour.
10 The resin was collected by allowing it to settle and decanting off the conditioned
media. The decanted media was filtered through a fritted disc filter to recover
any unsettled resin. The settled resin and that recovered by filtration was
resuspended and poured into a 2.5 cm diameter column and washed with three
column volumes of 0.15 M NaCl, 20 mM MES, pH 6.0 (A buffer). Protein was
15 eluted with a 300 ml gradient from A buffer to 1.0 M NaCl, 20 mM MES, pH 6.0
(B buffer), at a flow rate of 0.2 column volumes/minute with absorbance
monitored at 280 nm. Fractions containing 1.1 column volumes were collected.
The presence of GDNF in the fractions was detected by Western blotting analysis.
Fractions containing GDNF were pooled for further purification. GDNF eluted
20 between 0.3 and 0.6 M NaCl.

Step 2. HPLC C4 Chromatography:

The pool from Step 1 was made 0.1% (v/v) trifluoroacetic acid (TFA),
vacuum filtered through a 0.45 micron filter, and applied to a Vydac C4 column
(0.46 X 25 cm) conditioned with aqueous 10% acetonitrile, 0.1% TFA (A
25 buffer). Protein was eluted with a 2%/minute linear gradient over 50 minutes
from A buffer to aqueous 90% acetonitrile, 0.1% TFA (B buffer) with
absorbance measured at 280 nm. One milliliter fractions were collected, and the
presence of GDNF was detected by Western blotting analysis. GDNF was eluted
between 45% and 55% acetonitrile. Fractions were taken to dryness in vacuum.

Step 3. High performance S Chromatography:

Fractions containing GDNF from Step 2 were resolubilized in one
milliliter of 0.15 M NaCl, 10 mM Tris, pH 8.0, and applied to a 0.75 X 7.5 cm
TSK-Gel 5WP high performance S column (Toso Haas). A linear gradient of
35 0.4%/minute was run from 0.15 M NaCl, 10 mM Tris, pH 8.0 (A buffer) to
1.0 M NaCl, 10 mM Tris, pH 8.0 (B buffer) over 50 minutes at a flow rate of

1 ml/min. One minute fractions were collected with absorbance measured at 280 nm. At 35% B buffer, the gradient was changed to 6.5%/minute over 10 minutes. Western blot analysis of the fractions showed four major GDNF components. Three of the components eluted during the 0.4%/minute gradient and the fourth was eluted during the 6.5%/minute gradient. Appropriate pools were made of similar components and submitted for sequencing. Sequencing analyses identified an approximately 29-36 kD pool as [Arg³²-Ile¹³⁴] truncated GDNF protein. The component at approximately 38-40 kD was identified as an [Arg³²-Ile¹³⁴] truncated GDNF/mature GDNF heterodimer. Finally, the approximately 41-44 kD component which was isolated during the latter portion of the gradient was identified by sequencing as the mature GDNF homodimer.

Example 2

Mature Human GDNF Produced in *E. coli*

The bacterial expression of mature human GDNF may be achieved in accordance with the process described in Lin et al. (U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254); the disclosures of which are hereby incorporated by reference. Based upon the description of the present invention, those of ordinary skill in the art will appreciate that a variety of materials and methods may readily be used or adapted for suitable expression in *E. coli* and other bacteria. For example, alternate polynucleotides, such as those depicted in Figures 1, 3 and 4, may be used in the expression process.

Refolding and Purification of Expressed Mature GDNF

The transformed cells were processed at 5°C (unless otherwise noted) as follows: cell paste (30 gm) was suspended into a final volume of 200 milliliters using 25 mM Tris, pH 8.5 containing 5 mM EDTA, to yield a final cell slurry of 15% (w/v). The cells were thoroughly dispersed using a Biospec hand-held low shear homogenizer. The slurry was passed twice through a microfluidizer at 14,500 psi to break the cells and release inclusion bodies. The resulting

homogenate was then centrifuged for 30 minutes at 16,000xg. The pellet of inclusion bodies resulting from the centrifugation was washed by resuspension in chilled water to a final volume of 240 milliliters using the Biospec homogenizer, as before, to form a slurry. A sample of this slurry was kept for HPLC analysis of the GDNF expression level. The remaining slurry was centrifuged for 30 minutes at 16,000xg. The supernatant was discarded, and a small amount of cold water was added to the centrifuge bottle and gently swirled to remove the loosely formed membrane layer on top of the inclusion bodies pellet. The pellet was resuspended with the Biospec homogenizer using a sufficient volume of cold water to yield a concentration of 2 mg/ml of GDNF. The inclusion bodies were then solubilized by mixing the final inclusion bodies suspension (25 ml) and 8M guanidine HCl (25 ml) containing 180 mM cysteine HCl, and 50 mM Tris HCl, pH 8.7. The solubilization mixture was stirred at 25°C for 60 to 90 minutes after which it was poured, with mixing, into 2 M urea (450 ml, at 5°C) containing 20 mM Tris HCl, pH 8.75 and 0.2 M guanidine HCl. This refold mixture was slowly stirred at 5°C for 72 hours.

The refolded GDNF was partially purified as follows: 20 mM sodium acetate buffer (250 ml, pH 5) at 5°C was added with rapid stirring to the refold mixture, and the pH was adjusted to 5 with glacial acetic acid. The resulting precipitate was removed by centrifugation at 13,600xg for 45 minutes at 5°C. The supernatant from this centrifugation was used as the load solution for the next purification step involving cation exchange chromatography with SP-big bead resin (Pharmacia). The column was operated at 5°C using 20 mM sodium acetate (pH 5) as the equilibration, rinsing, and elution buffer system. A bed of resin (5 ml) was sanitized with 5 column volumes (CV) of 0.2N NaOH and then equilibrated with the acetate buffer (5 CV). The load solution (190 ml) was applied to the column at 0.5 CV/minute followed by a 10 CV rinse with acetate buffer at the same flow rate. The GDNF was then eluted off the resin with a 20 CV linear gradient of NaCl from 0.3 M to 0.9M in the acetate buffer at a flow rate of 0.1 CV/minute. The column eluate was monitored by absorbance at 280 nm and collected as fractions which were assayed by SDS-PAGE. The fractions containing GDNF were pooled from the front of the GDNF peak at 10% peak height through to the back of the peak to 10% peak height. The protein in this pool was entirely GDNF and, depending upon the production strain used, contained 32% to 12% contamination as modified GDNF forms. The pool was then dialyzed against PBS or other formulation buffers and, in some cases, concentrated by ultrafiltration

to 25 mg/ml. Both wild type and analogue forms of GDNF purified by this procedure were characterized by reverse phase HPLC, cation exchange HPLC, mass spectrometry, and endotoxin levels in order to compare purities of the preparations in relation to the corresponding production strains.

5

Example 3

Recombinant Production of Truncated GDNF in *E. coli*

10 Exemplary truncated GDNF proteins were produced substantially in accordance with the techniques described in Lin et al. (U.S. Patent Application No. 08/182,183 filed May 23, 1994, *supra*). Alternative bacterial expression materials and methods, as described above, may also be used. The *E. coli*-expressed truncated GDNF proteins included the [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], and [Gly³³-Ile¹³⁴] truncated GDNF proteins as depicted in Figures 5, 6 and 7, respectively. The polynucleotides encoding these exemplary truncated GDNF proteins were constructed as depicted in Figures 5, 6 and 7, but corresponding polynucleotides such as those depicted in Figures 1, 3 and 4 may also be used. The polynucleotides were constructed by standard PCR procedures as described in *PCR Technology, Principles and Applications for DNA Amplification*, Henry A. Erlich, ed., Stockton Press, NY, 1989 (Chapter 6, Using PCR to Engineer DNA) the disclosure of which is hereby incorporated by reference.

20
25

Example 4

Bioassay for Dopaminergic Neuron Neurotrophic Activity

The *E. coli*-expressed [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins of Example 3 and the CHO-derived [Arg³²-Ile¹³⁴] truncated GDNF protein of Example 1 were assessed on a qualitative basis for their ability to promote dopamine uptake by substantia nigra dopaminergic neurons.

35 Materials

The following materials are used in an assay to assess the survival of dopaminergic neurons in the presence of truncated GDNF proteins:

Cell Culture Media

High glucose Dulbecco's Modified Eagle's Medium (DMEM; catalog #11965-092), Ham's F12 medium (F12; #11765-021), Leibovitz's L15 medium without sodium bicarbonate (#41300-039), B27 medium supplement (#17504-010), penicillin/streptomycin (#15070-014), L-glutamine (#25030-016), Dulbecco's phosphate-buffered saline (D-PBS; #14190-052), Hank's balanced salt solution with calcium and magnesium salts (HBSS; #24020-026), N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES; #15630-015), mouse laminin (#23017-015), and bovine serum albumin fraction V (#110-18-017) were all from GIBCO, Grand Island, NY. Heat-inactivated horse serum was from HyClone, Logan, Utah. Conalbumin (C-7786), poly-L-ornithine hydrobromide (P-3655), bovine insulin (I-5500), human transferrin (T-2252), putrescine (P-6024), progesterone (P-6149), sodium selenite (S-9133), metrizamide (M-3383) were all from Sigma Chemical Company, Saint-Louis, MO. Papain, deoxyribonuclease I (DNAase) and ovalbumin (Papain dissociation system) were from Worthington Biochemicals, Freehold, NJ.

Falcon sterile 96-well microplates (#3072), tissue culture plastic ware and polypropylene centrifuge tubes were from Becton-Dickinson, Oxnard, CA. Nunc Lab-Tek tissue culture chamber coverglasses (#136439) were from Baxter, Irvine, CA; 20 μ m (#460) nylon mesh was from Tetko, Elmsford, NY; and 4" dissecting forceps and 4" dissecting scissors were from Roboz Surgical, Washington, DC.

Antibodies and Related Reagents

Polyclonal rabbit anti-tyrosine hydroxylase antibodies (TE101) were from Eugene Tech, Ridgefield Park, NJ; polyclonal rabbit anti-neuronal-specific enolase antibodies (NSE, AB951) were from Chemicon, Temecula, CA; and biotinylated goat anti-rabbit IgG and peroxidase-conjugated avidin/biotin complex (ABC Elite; Vectastain kit PK-6100) were from Vector Laboratories, Burlingame, CA. 3',3'-diaminobenzidine was from Cappel Laboratories, West Chester, PA. Superblock blocking buffer in PBS (#37515) was from Pierce Chemical Company, Rockford, IL. Triton X-100 (X100), Nonidet P-40 (N6507) and hydrogen peroxide (30%, v/v; H1009) were from Sigma. GBR-12909 dopamine uptake inhibitor (D-052) was from RBI, Natick, MA. ^3H -

dopamine (tritiated dopamine, NE-131; 21 Ci/mmol) was from New England Nuclear, Boston, MA. Optiphase Supermix scintillation cocktail was from Wallac, Turku, Finland. White ViewPlate-96 microplates (#6005182) were from Packard Instruments Corporation, Meriden, CT. All other reagents were

5 obtained from Sigma Chemical Company, unless otherwise specified.

Preparation of Media

The basal medium was prepared as a 1:1 mixture of DMEM and F12 medium, and was supplemented with B27 medium supplement added as a 50-fold

10 concentrated stock solution. L-glutamine was added at a final concentration of about 2 mM, penicillin at about 100 IU/l, and streptomycin at about 100 mg/l. Heat-inactivated horse serum was added to a final concentration of about 15 percent. After mixing, the pH was adjusted to about 7.3, and the media were kept

15 at 4°C. The media were prepared fresh just before use in order to minimize inter-experimental variations. Plastic pipettes and containers were used throughout to minimize protein adsorption.

Culture Substratum

To encourage optimal attachment of substratum neurons and neurite

20 outgrowth, microtiter plate surfaces (the culture substratum) were modified by sequential coatings with poly-L-ornithine and laminin, as follows. The plate surfaces were completely covered with a 0.1 mg/ml sterile solution of poly-L-ornithine in 0.1 M boric acid (pH 8.4) for at least one hour at room temperature, followed by a sterile wash with Super-Q water. The water wash

25 was then aspirated, and a 1.0 µg/ml solution of mouse laminin in PBS was added and incubated at 37°C for two hours. These procedures were conducted just before using the plates in order to ensure reproducibility of the results.

Preparation of Embryonic Rat Substantia Nigra Cultures

Embryonic rat brains were used as the source of substantia nigra

30 dopaminergic neurons. Timed-pregnant Sprague-Dawley rats at embryonic day 15 were used. A maximum of 36 embryos (about three litters) were processed per experiment. The pregnant rats were killed by exposure to CO₂, their abdominal cavities opened with dissecting scissors, and the fetuses were

35 removed from the uteri. Fetal brains were then dissected and cleaned of blood and meninges, and the ventral tegmental area containing the substantia nigra was

dissected using well-defined anatomical landmarks (Altman and Bayer, *Atlas of Prenatal Rat Brain Development*, CRC Press, Boca Raton, FL, 1995). The tissues were collected in ice-cold D-PBS, transferred into 10 milliliters of dissociation medium (120 units papain and 2000 units DNAase in HBSS) and then incubated for 45 minutes at about 37°C on a rotary platform shaker at about 200 rpm. The cells were then dispersed by trituration through fire-polished Pasteur pipettes, sieved through a 20 µm Nitex mesh to discard undissociated tissue, and centrifuged for five minutes at 200xg using an IEC clinical centrifuge. The resulting cell pellet was resuspended into HBSS containing ovalbumin and about 500 units DNAase, layered on top of a 4% ovalbumin solution (in HBSS) and centrifuged for about 10 minutes at 500xg. The final pellet was resuspended in complete culture medium (see above), adjusted to about 28,000 cells/ml, and seeded in aliquots (90 µl) into the 6 mm-wells of the 96-well microplates previously coated with polyornithine and laminin. Attachment of cells occurred rapidly, and the plating efficiency was about 75 percent.

Immunohistochemistry of Dopaminergic Neurons

An indirect immunoperoxidase method described by Louis et al. (*J. Pharmacol. Exp. Therap.*, 262:1274-1283, 1992; *Science*, 259:689-692, 1993) was used with slight modifications, as follows, to characterize the dopaminergic neurons in cultures of substantia nigra. Cultures were fixed for about 30 minutes at room temperature with 4% paraformaldehyde in D-PBS, pH 7.4, followed by three washes in D-PBS (200 µl per 6-mm well). The fixed cultures were then incubated in Superblock blocking buffer in PBS, containing 1% NP-40 to increase the penetration of the antibodies. The primary rabbit anti-tyrosine hydroxylase antibodies were then applied at a dilution of about 1:2000 in the same buffer and incubated for one hour at 37°C on a rotary shaker. After three washes with D-PBS, the bound antibodies were detected using goat-anti-rabbit biotinylated IgG at about a 1:500 dilution; these secondary antibodies were incubated with the cells for about one hour at 37°C. The cells were then washed three times with D-PBS, and the secondary antibodies were detected with avidin-biotin-peroxidase complex diluted at 1:500, and the cells were incubated for about 45 minutes at 37°C. After three more washes with D-PBS, the cultures were reacted for 5-20 minutes in a

solution of 0.1 M Tris-HCl, pH 7.4, containing 0.04% 3',3'-diaminobenzidine-(HCl)4, 0.06 percent NiCl₂ and 0.02 percent hydrogen peroxide.

Determining Neuronal Survival

- 5 Substantia nigra cultures were fixed and processed for immunostaining as described above, and then examined with bright-light optics at 200X magnification. The number of neurons stained for tyrosine hydroxylase was counted in the entire 6-mm well of the 96-well microplates. Viable neurons were characterized as having a regularly-shaped cell body, with a major axon-
10 like process and several dendrite-like processes. Neurons showing signs of degeneration, such as having irregular, vacuolated perikarya or fragmented neurites, were excluded from the counts (most of the degenerating neurons, however, detached from the culture substratum). Dopaminergic neuron cell numbers were expressed either as TH-positive neurons/6-mm well or as the
15 fold-change relative to control dopaminergic neuron density.

Determining Dopamine Uptake

- Dopamine uptake was determined in cultures of 15-day-old embryonic rat substantia nigra neurons that had been established in white ViewPlate-96
20 microplates. The cultures were washed with pre-warmed uptake buffer (about 100 µl) which consists of a modified Krebs-Ringer solution, pH 7.4, containing about 120 mM NaCl, 4.7 mM KCl, 1.8 mM CaCl₂, 1.2 mM MgSO₄, 32 mM NaHPO₄, 1.3 mM EDTA, and 5.6 mM D-glucose. The uptake buffer also contained 1 mM ascorbic acid and 50 µM pargyline to prevent the oxidation of dopamine.
25 The cells were then preincubated at 37°C for about 10 minutes in uptake buffer. Tritiated dopamine (³H-DA, 21 Ci/mmol) was then added to the substantia nigra cultures at a concentration of about 50 nM in 75 µl of uptake buffer, and the cultures were incubated for about 60 minutes at 37°C. Non-specific dopamine uptake was determined by incubating the cultures with uptake buffer containing
30 the dopamine uptake inhibitor GBR-12909 (1 µM). Non-specific uptake represented less than about one percent of total uptake. The uptake assays were arrested by aspiration of the incubation medium followed by three rapid washes with ice-cold uptake buffer (about 120 µl). The cells were then lysed by addition of Optiphase Supermix scintillation cocktail (200 µl), and radioactivity
35 was determined by scintillation spectrometry using a Wallac MicrobetaPlus 96-well microplate counter (i.e., dopamine uptake is analyzed by scintillation

counting of the retained tritium in the cultures). The results are expressed either as dpm/6-mm well or as the fold-change relative to control cultures.

Assays

5 Dopaminergic Neuron Survival and Morphological Development

Cultures of embryonic day 15 (E15) rat substantia nigra enriched in dopaminergic neurons were used to demonstrate the effect of truncated GDNF proteins on the survival of dopaminergic neurons. The cultures were grown in polyornithine- and laminin-coated 96-well microplates for up to six days alone
10 or in the presence of various concentrations (ranging from about 1 pg/ml to about 10 ng/ml) of the following proteins: *E. coli*-expressed mature hGDNF; *E. coli*-expressed [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins; CHO cell-expressed mature hGDNF; and CHO cell-derived [Arg³²-Ile¹³⁴] truncated GDNF protein. The culture
15 medium consisted of DMEM/F12 supplemented with 15% heat-inactivated horse serum (E15 cultures) or 2.5% heat-inactivated horse serum, D-glucose, HEPES, insulin and transferrin (P6 cultures). Immunostaining for tyrosine hydroxylase (TH), the rate-limiting enzyme in dopamine biosynthesis, was used as a marker for dopaminergic neurons. Since noradrenergic neurons in the
20 rhombencephalon also stain positive for TH, great care was taken to dissect an area restricted to the ventral tegmentum of the mesencephalon and to avoid the more caudal regions containing the noradrenergic cell bodies. After six days, the E15 cultures typically consisted of about 70% neurons as identified by neuronal specific enolase immunostaining (described above) and 30% non-neuronal cells
25 (which had a flattened, phase-dark appearance); dopaminergic neurons represented about 10-15% of the neuron population.

After six days, the cultures were fixed with paraformaldehyde and immunostained for tyrosine hydroxylase, a marker that identifies dopaminergic neurons in these cultures. All the tyrosine-hydroxylase-positive neurons
30 present in a 6-mm well were counted under brightfield optics. Three to six different wells were analyzed for each experimental condition. The results were expressed as the percentages of the number of tyrosine-hydroxylase-positive neurons found in control cultures.

Cultures of E15 substantia nigra treated with 1.0 ng/ml of GDNF, CHO
35 cell-expressed GDNF or *E. coli*-expressed GDNF, contained about 38% and 27% more TH-immunoreactive neurons, respectively, than untreated control

cultures, indicating that both GDNF species promote the survival of dopaminergic neurons. Cultures of E15 substantia nigra treated with 1.0 ng/ml of truncated GDNF protein showed a similar increase in the number of TH-positive neurons in cultures after six days *in vitro* : 42% for CHO cell-derived [Arg³²-Ile¹³⁴] truncated GDNF protein; and 26% and 17% for *E. coli*-expressed [Arg³²-Ile¹³⁴] and [Gly³³-Ile¹³⁴] truncated GDNF proteins, respectively.

Comparison of control cultures and cultures treated with mature and truncated GDNF proteins also revealed pronounced effects of all the GDNF proteins on the morphological differentiation of dopaminergic neurons. The effects of the [Arg³²-Ile¹³⁴] and [Gly³³-Ile¹³⁴] truncated GDNF proteins were identical to their respective mature GDNF protein counterparts. TH-immunoreactive neurons in all GDNF-treated cultures possessed significantly more complex and extensive neuritic arborization, as well as a higher degree of neurite branching and an overall larger soma size, than did TH-positive neurons in control cultures.

Dopamine Uptake

Dopamine uptake measures the number and activity of high affinity dopamine reuptake transporter sites and reflects the functional differentiation of dopaminergic neurons. Dopamine uptake was measured in cultures of E15 rat substantia nigra after six days *in vitro* either with or without mature GDNF or truncated GDNF proteins. In these cultures, dopamine uptake had the pharmacological profile characteristic of dopaminergic neurons, i.e., it was nearly completely blocked (more than 98 percent) by 1.0 μ M GBR-12909, a dopamine transporter inhibitor specific for dopaminergic neurons (ID₅₀ = 20 nM). This indicates that dopamine uptake measurements do not reflect the presence of contaminating noradrenergic neurons, which can take up dopamine through norepinephrine transporters but are not sensitive to GBR-12909 inhibition. The effects of CHO-cell-expressed mature GDNF and the CHO-derived [Arg³²-Ile¹³⁴] truncated GDNF protein were identical: about 65% increase, with an ED₅₀ of about 20 pg/ml. *E. coli*-expressed [Pro²³-Lys³⁷ Δ Asn³⁷-Ile¹³⁴] truncated GDNF protein, as depicted in Figure 5, demonstrated a 65% increase, with an ED₅₀ of about 40 pg/ml. The effects on dopamine uptake of the *E. coli*-expressed mature protein and the *E. coli*-expressed [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins were the same: about 50% increases, with ED₅₀s of about 50 pg/ml.

These results indicate that the truncated GDNF proteins act as potent survival-promoting and differentiation-inducing factors for substantia nigra dopaminergic neurons. As such, they are envisioned to be particularly useful in the treatment of Parkinson's disease, a neurological disorder characterized by

5 decreased emotional acuity, slowing of both voluntary and involuntary muscle movement, muscular rigidity, and tremor. Such symptoms are attributed to the progressive degeneration of dopamine-producing neurons located in the substantia nigra. Degeneration of these neurons ("dopaminergic neurons") results in a decrease of dopamine in an adjacent region of the brain called the


10 striatum.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209

CLAIMS

What is claimed is:

- 5 1. A truncated glial cell line-derived neurotrophic factor (GDNF) protein product having an amino acid sequence



wherein

- [Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through
10 Cys¹³³ as depicted in Figure 1 (SEQ ID NO 2);

Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

15

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GQRGKNRG (SEQ ID NO:7)
RGQRGKNRG (SEQ ID NO:8)
RRGQRGKNRG (SEQ ID NO:9)
G RRGQRGKNRG (SEQ ID NO:10)
KG RRGQRGKNRG (SEQ ID NO:11)
GKG RRGQRGKNRG (SEQ ID NO:12)
RGKG RRGQRGKNRG (SEQ ID NO:13)
SRGKG RRGQRGKNRG (SEQ ID NO:14)
NSRGKG RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG RRGQRGKNRG (SEQ ID NO:17)
NPENSRGKG RRGQRGKNRG (SEQ ID NO:18)
ANPENSRGKG RRGQRGKNRG (SEQ ID NO:19)
A ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)

AA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:21)
AAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:22)
QAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:23)
RQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:24)
NRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:25)
RNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:26)
ERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:27)
RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:28)
RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:29)
P RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:30)
LP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:31)
VLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:32)
AVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:33)
MAVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:34)
QMAVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:35)
KQMAVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:36)
DKQMAVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:37) and
PDKQMAVLP RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:38)

and addition, substitution and internal deletion variants and derivatives thereof.

2. A truncated GDNF protein product according to Claim 1, wherein X =
 5 RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24) or a variant thereof.

3. A truncated GDNF protein product according to Claim 1, wherein X =
 NPENSRGKG RRGQRGKNRG (SEQ ID NO:18) or a variant thereof.

10 4. A truncated GDNF protein product according to Claim 1, wherein X =
 PENSRGKG RRGQRGKNRG (SEQ ID NO:17) or a variant thereof.

5. A truncated GDNF protein product according to Claim 1, wherein X =
 SRGKG RRGQRGKNRG (SEQ ID NO:14) or a variant thereof.

15

6. A truncated GDNF protein product according to Claim 1, wherein X =
 RRGQRGKNRG (SEQ ID NO:8) or a variant thereof.

7. A truncated GDNF protein product according to Claim 1, wherein X = GQRGKNRG (SEQ ID NO:7) or a variant thereof.
8. A truncated GDNF protein product according to Claim 1, wherein X =
5 KNRG (SEQ ID NO:3) or a variant thereof.
9. A truncated GDNF protein product according to Claim 1, wherein X = NRG or a variant thereof.
- 10 10. A truncated GDNF protein product according to Claims 1 through 9, wherein said amino acid sequence is glycosylated.
11. A truncated GDNF protein product according to Claims 1 through 9, wherein said amino acid sequence is nonglycosylated.
- 15 12. A truncated GDNF protein product according to Claim 1, wherein said derivative is an X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence conjugated to a water soluble polymer.
- 20 13. A polynucleotide encoding a truncated GDNF protein according to Claim 1.
14. A polynucleotide according to Claim 13, comprising a portion of the sequence as set forth in Figure 1.
- 25 15. A polynucleotide according to Claim 13, comprising a portion of the sequence as set forth in Figure 3.
16. A polynucleotide according to Claim 13, comprising a portion of the sequence as set forth in Figure 4.
- 30 17. A polynucleotide according to Claim 13, comprising the sequence as set forth in Figure 5.
18. A polynucleotide according to Claim 13, comprising the sequence as set
35 forth in Figure 6.

19. A polynucleotide according to Claim 13, comprising the sequence as set forth in Figure 7.

20. A vector, comprising a polynucleotide of Claim 13 operatively linked to an expression control sequence.

21. A prokaryotic or eukaryotic host cell transformed or transfected with a polynucleotide of Claim 13.

22. A method for producing a truncated GDNF protein, comprising growing host cells of Claim 21 in a suitable nutrient medium and, optionally, isolating said truncated GDNF from said cells or said nutrient medium.

23. A method for producing a truncated GDNF protein according to Claim 22, wherein said host cells are *E. coli*.

24. A method for producing a truncated GDNF protein according to Claim 22, wherein said host cells are Chinese hamster ovary cells.

25. A method for the production of a truncated glial cell line-derived neurotrophic factor (GDNF) protein, comprising the steps of:

- (a) culturing a prokaryotic or eukaryotic host cell transformed or transfected with a vector of Claim 20;
- (b) maintaining said host cell under conditions allowing the expression of truncated GDNF protein by said host cell; and
- (c) optionally isolating the truncated GDNF protein expressed by said host cell.

26. A truncated GDNF protein which is the recombinant expression product of a prokaryotic or eukaryotic host cell containing an exogenous polynucleotide of Claim 13.

27. A pharmaceutical composition comprising a truncated GDNF protein product according to Claim 1 in association with a pharmaceutically acceptable vehicle.

28. A pharmaceutical composition comprising a truncated GDNF protein, produced in accordance with the method of Claim 22 in association with a pharmaceutically acceptable vehicle.

5 29. A pharmaceutical composition comprising a truncated GDNF protein, produced in accordance with the method of Claim 25 in association with a pharmaceutically acceptable vehicle.

10 30. A method of treating Parkinson's Disease comprising administering to a patient the pharmaceutical composition of Claim 27.

15 31. A method of treating Parkinson's Disease comprising administering to a patient a polynucleotide sequence of Claim 13 to provide *in vivo* production of said truncated GDNF protein.

32. A method of treating Parkinson's Disease comprising implanting in a patient a cell transformed with a polynucleotide sequence of Claim 13 to provide *in vivo* production of said truncated GDNF protein.

20 33. A glial cell line-derived neurotrophic factor (GDNF) composition, comprising a mature GDNF protein and one or more truncated GDNF proteins, wherein said mature GDNF protein has a molecular weight of approximately 44 kDa, and wherein said truncated GDNF protein(s) has a molecular weight of approximately 29 to 40 kDa.

25 34. A GDNF composition according to Claim 33, comprising at least two species of truncated GDNF protein wherein a first species has a molecular weight of approximately 36 kDa and a second species has a molecular weight of approximately 40 kDa.

30 35. A GDNF composition according to Claim 34, wherein said second truncated GDNF species having a molecular weight of approximately 40 kDa is a heterodimer of a mature GDNF monomer having a molecular weight of approximately 22 kDa and a truncated GDNF monomer having a molecular weight of approximately 18 kDa.

35

36. A truncated GDNF protein isolated from the GDNF composition of Claim 33 and having a molecular weight of approximately 29 to 40 kDa.

37. A truncated GDNF protein isolated from the GDNF composition of Claim 33 and having a molecular weight of approximately 29 to 36 kDa.

38. A truncated GDNF protein derived from a mature GDNF protein expressed by a recombinantly modified bacterial or mammalian cell, said truncated GDNF protein having an amino acid sequence

10 X-[Cys⁴¹-Cys¹³³]-Y

wherein

[Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO 2);

15 Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

X represents an amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GQRGKNRG (SEQ ID NO:7)
RGQRGKNRG (SEQ ID NO:8)
RRGQRGKNRG (SEQ ID NO:9)
G RRGQRGKNRG (SEQ ID NO:10)
KG RRGQRGKNRG (SEQ ID NO:11)
GKG RRGQRGKNRG (SEQ ID NO:12)
RGKG RRGQRGKNRG (SEQ ID NO:13)
SRGKG RRGQRGKNRG (SEQ ID NO:14)
NSRGKG RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG RRGQRGKNRG (SEQ ID NO:17)

	NPENSRGKG	RRQRGKNRG	(SEQ ID NO:18)
	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:19)
A	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:20)
AA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:21)
AAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:22)
QAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:23)
RQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:24)
NRQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:25)
RNRQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:26)
ERNRQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:27)
RRERNRQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:28)
RRERNRQAAA	ANPENSRGKG	RRQRGKNRG	(SEQ ID NO:29)
P	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:30)
LP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:31)
VLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:32)
AVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:33)
MAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:34)
QMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:35)
KQMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:36)
DKQMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:37) and
PDKQMAVLP	RRERNRQAAA	ANPENSRGKG	RRQRGKNRG (SEQ ID NO:38)

and addition, substitution and internal deletion variants thereof.

39. A truncated GDNF protein according to Claim 38, wherein X is selected
 5 from the group consisting of:

G
 RG
 NRG
 KNRG (SEQ ID NO:3)
 GKNRG (SEQ ID NO:4)
 RGNRG (SEQ ID NO:5)
 QRGKNRG (SEQ ID NO:6)
 GQRGKNRG (SEQ ID NO:7)
 RGQRGKNRG (SEQ ID NO:8) and
 RRQRGKNRG (SEQ ID NO:9)

and variants thereof.

40. A truncated GDNF protein according to Claim 38, wherein mature GDNF protein is expressed by a recombinantly modified bacterial cell and said truncated GDNF protein is produced *in vitro* or *in vivo*.

41. A method of preparing a pharmaceutical composition wherein a therapeutically effective amount of a truncated GDNF protein product according to Claim 1 is mixed with one or more pharmaceutically acceptable vehicles.

10

42. The use of a truncated GDNF protein product according to Claim 1 for treating damage to the nervous system caused by disease or injury.

43. The use of a truncated GDNF protein product according to Claim 42 for treating Parkinson's disease.

15

44. The use of a truncated GDNF protein product according to Claim 1 for the preparation of a pharmaceutical composition for treating damage to the nervous system caused by disease or injury.

20


TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR**ABSTRACT**

5 Disclosed are novel proteins, referred to as truncated glial cell line-derived neurotrophic factor (truncated GDNF) proteins, that promote dopamine uptake by dopaminergic cells and promote the survival of nerve cells. Also disclosed are processes for obtaining the truncated GDNF proteins by recombinant genetic engineering techniques.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
22

FIG. 2



1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100

[illegible][illegible]

FIG.3B

S
u
n
I

241 CCTGCGACGCTGCTGAAACCACGTACGACAAAATCCTGAAAAACCTGTCCCGTAACCGTC 300

H i n d I P v u I

301 GTCTGGTTTCCGACAAAGTTGGTCAAGCTTGTGCGTCCGATCGCTTTCGACGACGACC 360

B a m H I

361 TGTCC TTCCTGGACGACAACCTGGTTTACCACATCCTGCGTAAACACTCCGCTAAGCGTT 420

421 GCGGTTGCATCTAAGGATCC 440

FIG.4

metGDNF Degenerate DNA Sequence

```

                                N
                                d
                                e
                                I
                                CATATGAGCCCGGACAAACAG
1  -----+-----+-----+-----+-----+ 60
                                MetSer

ATGGCAGTACTTCCACGTCGTGAACGTAATCGCCAGGCAGCAGCTGCAAACCCGAAAAAC
61 -----+-----+-----+-----+-----+ 120

TCCCGTGGTAAAGGTCGCCGTGGCCAGCGCGGCAAAAACCGTGGTGTGTCTGACTGCA
121 -----+-----+-----+-----+-----+ 180

                                P
                                s
                                h
                                A
                                I
181 ATCCACCTGAACGTTACTGACCTGGGTCTGGGCTACGAAACCAAGAAGAACTGATCTTC
-----+-----+-----+-----+-----+ 240

                                P
                                s
                                t
                                I
241 CGTACTGCAGCGGCTCTTGCGACGCAGCTGAAACCACTTACGACAAAATCCTGAAAAAC
-----+-----+-----+-----+-----+ 300

                                                                P
                                                                v
                                                                u
                                                                I
301 CTGTCCCCTAACC GCCGTCTTGTAAGCGACAAAGTAGGT CAGGCATGCTGCCGTC CGATC
-----+-----+-----+-----+-----+ 360

B
s
m
I
361 GCATTCGACGATGACCTGAGCTTCTGGATGACAACCTGGTTTACCACATCCTGCGTAA
-----+-----+-----+-----+-----+ 420

                                B
                                a
                                m
                                H
                                I
421 CACTCCGCTAAACGCTGCGGTGTCATCTAAGGATCC
-----+-----+-----+-----+ 456

```

FIG.5

[Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] Truncated GDNF Protein

ATGTCCCCAGAAAATTCTCGTGGTAAAGGTCGTCGTCGGTCAGCGTGGTAATAACCGCGGT
21 -----+-----+-----+-----+-----+-----+ 80
M S P E N S R G K G R R G Q R G N N R G
TGC GTTCTGACCGCTATCCACCTGAACGTTACCGACCTGGGTCTCGGTTACGAAACCAAA
81 -----+-----+-----+-----+-----+-----+ 140
C V L T A I H L N V T D L G L G Y E T K
GAAGAATTAATCTTCCGTTACTGCTCCGTTTCTGCGACGCTGCTGAAACCACGTACGAC
141 -----+-----+-----+-----+-----+-----+ 200
E E L I F R Y C S G S C D A A E T T Y D
AAAATCCTGAAAAACCTGTCCCGTAACCGTCGTCTGGTTTCCGACAAAGTTGGTCAAGCT
201 -----+-----+-----+-----+-----+-----+ 260
K I L K N L S R N R R L V S D K V G Q A
TGCTGCCGTCGGATCGCTTTCGACGACGACCTGTCCTTCCTGGACGACAACCTGGTTTAC
261 -----+-----+-----+-----+-----+-----+ 320
C C R P I A F D D D L S F L D D N L V Y
CACATCCTGCGTAAACACTCCGCTAAGCGTTGCGGTTGCATCTAA
321 -----+-----+-----+-----+-----+-----+
H I L R K H S A K R C G C I *

FIG.6

[Arg³²-Ile¹³⁴] Truncated GDNF Protein

ATGCGTGGTCAACGTGGTAAAAACCGCGTTGCGTTCTGACTGCAATCCACCTGAACGTT
41 -----+-----+-----+-----+-----+-----+ 100
M R G Q R G K N R G C V L T A I H L N V

ACTGACCTGGGTCTGGGCTACGAAACCAAGAAGAACTGATCTCCGCTACTGCAGCGGC
101 -----+-----+-----+-----+-----+-----+ 160
T D L G L G Y E T K E E L I F R Y C S G

TCTTGCAGCGCAGCTGAAACCACTTACGACAAAATCCTGAAAAACCTGTCCCGTAAACCGC
161 -----+-----+-----+-----+-----+-----+ 220
S C D A A E T T Y D K I L K N L S R N R

CGTCTGGTAAGCGACAAAGTAGGTCAGGCATGCTGCCGTCCGATCGCATTCGACGATGAC
221 -----+-----+-----+-----+-----+-----+ 280
R L V S D K V G Q A C C R P I A F D D D

CTGAGCTTCCTGGATGACAACTGGTTTACCACATCCTGCGTAAACACTCCGCTAAACGC
281 -----+-----+-----+-----+-----+-----+ 340
L S F L D D N L V Y H I L R K H S A K R

TGCGGTTGCATCTAA
341 -----+----- 355
C G C I *

FIG.7

[Gly³³-Ile¹³⁴] Truncated GDNF Protein

ATGGGTCAACGTGGTAAAAACCGTGGTTGTGTTCTGACTGCAATCCACCTGAACGTTACT
41 -----+-----+-----+-----+-----+-----+ 100
M G Q R G K N R G C V L T A I H L N V T

GACCTGGGTCTGGGCTACGAAACCAAGAAGAACTGATCTTCCGCTACTGCAGCGGCTCT
101 -----+-----+-----+-----+-----+-----+ 160
D L G L G Y E T K E E L I F R Y C S G S

TGCGACGCAGCTGAAACCACTTACGACAAAATCCTGAAAAACCTGTCCCGTAACCGCCGT
161 -----+-----+-----+-----+-----+-----+ 220
C D A A E T T Y D K I L K N L S R N R R

CTGGTAAGCGACAAAGTAGGTCAGGCATGCTGCCGTCCGATCGCATTCGACGATGACCTG
221 -----+-----+-----+-----+-----+-----+ 280
L V S D K V G Q A C C R P I A F D D D L

AGCTTCCTGGATGACAACCTGGTTTACCACATCCTGCGTAAACACTCCGCTAAACGCTGC
281 -----+-----+-----+-----+-----+-----+ 340
S F L D D N L V Y H I L R K H S A K R C

GGTTGCATCTAA
341 -----+--- 352
G C I *

FIG.8

Comparison of Protein Sequences

						50
	GDNF	MSPDKQMAVL	PRRERNRQAA	AANPENSRGK	GRRGQRGKNR	GCVLTAIHLN
-31	GDNFMRGQRGKNR	GCVLTAIHLN
-32	GDNFMGQRGKNR	GCVLTAIHLN
-22	GDNFMSPENSRGK	GRRGQRGNR	GCVLTAIHLN
						100
	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN	RRLVSDKVGQ
-31	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN	RRLVSDKVGQ
-32	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN	RRLVSDKVGQ
-22	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN	RRLVSDKVGQ
						135
	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI	
-31	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI	
-32	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI	
-22	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI	

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first, and sole inventor (if only one name is listed below) or a joint inventor (if plural names are listed below) of the invention entitled:

TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR

which is described and claimed in the specification which:

☒ is attached hereto.

☐ was filed on _____
as Application Serial No.: _____
and was amended on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

Power of Attorney: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Ron K. Levy, Registration No.: 31,539, Steven M. Odre, Registration No.: 29,094, and Daniel R. Curry, Registration No. 32,727, said attorney(s)/agent(s) to have in addition full power of revocation, including the power to revoke any power herein granted.

Please send all future correspondence to:

U.S. Patent Operations/DRC
M/S 10-1-B
AMGEN INC.
Amgen Center
1840 Dehavilland Drive
Thousand Oaks, California 91320-1789

Direct Telephone Calls To:

Daniel R. Curry
Attorney/Agent for Applicant(s)
Registration No.: 32,727
Phone: (805) 447-8102
Date: September 28, 1995

"Express Mail" mail labeling number: YB 530269575 US	Date of Deposit: September 28, 1995
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner of Patents, Washington, D. C. 20231 (Senders Signature Below)	
(typed name) Ronda G. Spahr	<i>Ronda G. Spahr</i>

DECLARATION AND POWER OF ATTORNEY (cont'd)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full Name of Sole

or First Inventor:

Shaw-Fen Sylvia Hu

Inventor's Signature:

*Shaw-Fen Sylvia Hu*Date: Sept 28, 1995

Residence:

986 Lynnmere Drive

Post Office Address:

Thousand Oaks, California 91360 U.S.A.

Citizenship:

U.S.A.

Full Name of Second

Joint Inventor, if Any:

Inventor's Signature:

Date: _____

Residence:

Post Office Address:

Citizenship:

Full Name of Third

Joint Inventor, if Any:

Inventor's Signature:

Date: _____

Residence:

Post Office Address:

Citizenship:

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Shaw-Fen Sylvia Hu
Serial No.: To Be Assigned
Filed: October 13, 2000
For: Truncated Glial Cell Line-Derived Neurotrophic Factor
Docket No.: A-357C

ATTORNEY'S STATEMENT PURSUANT TO 37 C.F.R. 1.821(e)


Assistant Commissioner for Patents
Washington, D. C. 20231

Sir:

The computer readable form in this Application, is identical with that filed in Application Number 08/535,681; filed September 28, 1995. The present application is a divisional of application Serial No. 08/535,681. In accordance with 37 CFR 1.821(e), please use the last-filed computer readable form filed in that application as the computer readable form for the instant application. It is understood that the Patent and Trademark Office will make the necessary change in application number and filing date for the computer readable form that will be used for the instant application. A paper copy of the Sequence Listing is included for incorporation into the specification.

In addition to (1) the paper copy included in the instant application and (2) the reference to the other application and computer readable form, the undersigned verifies that the content of the paper copy in the instant application is identical to the computer readable form filed in Application Number 08/535,681.

Respectfully submitted,



Daniel R. Curry
Attorney for Applicant
Registration No.: 32,7270
Phone: (805) 447-8102
Date October 13, 2000

Please send all future correspondence to:

U.S. Patent Operations/DRC
Dept. 4300, M/S 27-4-A
AMGEN INC.
One Amgen Center Drive
Thousand Oaks, California 91320-1799

EXPRESS MAIL CERTIFICATE

"Express Mail" mail labeling number: EL360688271US

Date of Deposit: October 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231

3. L. St. Andrew
Printed Name

3. L. St. Andrew
Signature

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Hu, Sylvia
- (ii) TITLE OF INVENTION: Truncated Glial Cell Line-Derived
Neurotrophic Factor
- (iii) NUMBER OF SEQUENCES: 50
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: AMGEN INC.
 - (B) STREET: 1840 DeHavilland Drive
 - (C) CITY: Thousand Oaks
 - (D) STATE: California
 - (E) COUNTRY: United States of America
 - (F) ZIP: 91320
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Curry, Daniel R.
 - (B) REGISTRATION NUMBER: 32,727
 - (C) REFERENCE/DOCKET NUMBER: A-357
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 805-447-8102
 - (B) TELEFAX: 805-499-8011
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 402 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 1..402

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCA CCA GAT AAA CAA ATG GCA GTG CTT CCT AGA AGA GAG CGG AAT CGG	48
Ser Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg	
1 5 10 15	
CAG GCT GCA GCT GCC AAC CCA GAG AAT TCC AGA GGA AAA GGT CGG AGA	96
Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg	
20 25 30	
GGC CAG AGG GGC AAA AAC CGG GGT TGT GTC TTA ACT GCA ATA CAT TTA	144
Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu	
35 40 45	
AAT GTC ACT GAC TTG GGT CTG GGC TAT GAA ACC AAG GAG GAA CTG ATT	192
Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile	
50 55 60	
TTT AGG TAC TGC AGC GGC TCT TGC GAT GCA GCT GAG ACA ACG TAC GAC	240
Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp	
65 70 75 80	
AAA ATA TTG AAA AAC TTA TCC AGA AAT AGA AGG CTG GTG AGT GAC AAA	288
Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys	
85 90 95	
GTA GGG CAG GCA TGT TGC AGA CCC ATC GCC TTT GAT GAT GAC CTG TCG	336
Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser	
100 105 110	
TTT TTA GAT GAT AAC CTG GTT TAC CAT ATT CTA AGA AAG CAT TCC GCT	384
Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala	
115 120 125	
AAA AGG TGT GGA TGT ATC	402
Lys Arg Cys Gly Cys Ile	
130	

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 134 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ser Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg
1 5 10 15
Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg
20 25 30
Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu
35 40 45

Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile
 50 55 60
 Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp
 65 70 75 80
 Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys
 85 90 95
 Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser
 100 105 110
 Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala
 115 120 125
 Lys Arg Cys Gly Cys Ile
 130

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Lys Asn Arg Gly
 1

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Lys Asn Arg Gly
 1 5

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
1 5 10 15

Gly

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
1 5 10 15

Arg Gly

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
1 5 10 15

Asn Arg Gly

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1 5 10 15

Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
 1 5 10 15

Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
 1 5 10 15

Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
 1 5 10 15
 Gln Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg
 1 5 10 15
 Gly Gln Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
 1 5 10 15
 Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly
1           5                      10                      15

Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
                20                      25

```

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```

Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys
1           5                      10                      15

Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
                20                      25

```

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

```

Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly
1           5                      10                      15

Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
                20                      25

```

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```

Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg
1           5                      10                      15

```

(2) INFORMATION FOR SEQ ID NO:29:

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
20 25 30

(2) INFORMATION FOR SEO ID NO:30:

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
20 25 30

(2) INFORMATION FOR SEQ ID NO:31:

- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu
1 5 10 15

Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro
 1 5 10 15

Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
 20 25 30

Gly

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn
 1 5 10 15

Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
 20 25 30

Arg Gly

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala
 1 5 10 15
 Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
 20 25 30
 Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala
 1 5 10 15
 Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 20 25 30
 Lys Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala
 1 5 10 15
 Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
 20 25 30
 Gly Lys Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

```

Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala
 1             5             10             15

Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
          20             25             30

Arg Gly Lys Asn Arg Gly
      35

```

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

```

Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln
 1             5             10             15

Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
          20             25             30

Gln Arg Gly Lys Asn Arg Gly
      35

```

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 417 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

```

CATATGTCTC CGGATAAACA AATGGCTGTT CTTCCACGTC GTGAACGTAA CCGTCAGGCG 60
GCCGCTGCTA ACCCGAGGAA TTCCCGTGGT AAAGGTCGTC GTGGTCAGCG TGGTAAAAAC 120
CGCGGTTGCG TTCTGACCGC TATCCACCTG AACGTTACCG ACCTGGGTCT CGGTTACGAA 180
ACCAAAGAAG AATTAATCTT CCGTTACTGC TCCGTTCTCT GCGACGCTGC TGAAACCACG 240

```

TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGTCGTC TGGTTTCCGA CAAAGTTGGT 300
 CAAGCTTGCT GCCGTCGAT CGCTTTCGAC GACGACCTGT CTTTCCTGGA CGACAACCTG 360
 GTTTACCACA TCCTGCGTAA ACACTCCGCT AAGCGTTGCG GTTGCACTTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 417 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CATATGAGCC CGGACAAACA GATGGCAGTA CTTCCACGTC GTGAACGTAA TCGCCAGGCA 60
 GCAGCTGCAA ACCCGGAAAA CTCCTGGTGGT AAAGGTCGCC GTGGCCAGCG CGGCAAAAC 120
 CGTGTTGTGT TTCTGACTGC AATCCACCTG AACGTTACTG ACCTGGGTCT GGGCTACGAA 180
 ACCAAGAAG AACTGATCTT CCGCTACTGC AGCGGCTCTT GCGACGCAGC TGAAACCACT 240
 TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGCCGTC TGGTAAGCGA CAAAGTAGGT 300
 CAGGCATGCT GCCGTCGAT CGCATTGAC GATGACCTGA GCTTCCTGGA TGACAACCTG 360
 GTTTACCACA TCCTGCGTAA ACACTCCGCT AAACGCTGCG GTTGCACTTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 345 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..342

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ATG TCC CCA GAA AAT TCT CGT GGT AAA GGT CGT CGT GGT CAG CGT GGT 48
 Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 135 140 145 150
 AAT AAC CGC GGT TGC GTT CTG ACC GCT ATC CAC CTG AAC GTT ACC GAC 96
 Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
 155 160 165

- 76 -

(2) INFORMATION FOR SEO ID NO:42:

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

[illegible]

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 315 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..312

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

```

ATG CGT GGT CAA CGT GGT AAA AAC CGC GGT TGC GTT CTG ACT GCA ATC   48
Met Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile
115                               120                               130

CAC CTG AAC GTT ACT GAC CTG GGT CTG GGC TAC GAA ACC AAA GAA GAA   96
His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
135                               140                               145

CTG ATC TTC CGC TAC TGC AGC GGC TCT TGC GAC GCA GCT GAA ACC ACT  144
Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
150                               155                               160

TAC GAC AAA ATC CTG AAA AAC CTG TCC CGT AAC CGC CGT CTG GTA AGC  192
Tyr Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser
165                               170                               175

GAC AAA GTA GGT CAG GCA TGC TGC CGT CCG ATC GCA TTC GAC GAT GAC  240
Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
180                               185                               190

CTG AGC TTC CTG GAT GAC AAC CTG GTT TAC CAC ATC CTG CGT AAA CAC  288
Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
195                               200                               205

TCC GCT AAA CGC TGC GGT TGC ATC TAA                               315
Ser Ala Lys Arg Gly Cys Ile
215

```

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

```

Met Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile
  1              5              10              15

```

A-357

- 78 -

```

His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
      20                      25                      30

Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
      35                      40                      45

Tyr Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser
      50                      55                      60

Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
      65                      70                      75                      80

Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
      85                      90                      95

Ser Ala Lys Arg Cys Gly Cys Ile
      100

```

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 312 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..309

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

```

ATG GGT CAA CGT GGT AAA AAC CGT GGT TGT GTT CTG ACT GCA ATC CAC      48
Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
105                      110                      115                      120

CTG AAC GTT ACT GAC CTG GGT CTG GGC TAC GAA ACC AAA GAA GAA CTG      96
Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
      125                      130                      135

ATC TTC CGC TAC TGC AGC GGC TCT TGC GAC GCA GCT GAA ACC ACT TAC      144
Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
      140                      145                      150

GAC AAA ATC CTG AAA AAC CTG TCC CGT AAC CGC CGT CTG GTA AGC GAC      192
Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
      155                      160                      165

AAA GTA GGT CAG GCA TGC TGC CGT CCG ATC GCA TTC GAC GAT GAC CTG      240
Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
      170                      175                      180

AGC TTC CTG GAT GAC AAC CTG GTT TAC CAC ATC CTG CGT AAA CAC TCC      288
Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
185                      190                      195                      200

```

GCT AAA CGC TGC GGT TGC ATC TAA
 Ala Lys Arg Cys Gly Cys Ile
 205

312

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 103 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1 5 10 15
 Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 20 25 30
 Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 35 40 45
 Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
 50 55 60
 Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
 65 70 75 80
 Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
 85 90 95
 Ala Lys Arg Cys Gly Cys Ile
 100

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 135 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Met Ser Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn
 1 5 10 15
 Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
 20 25 30
 Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 35 40 45

```

Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 50                               55                               60

Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 65                               70                               75                               80

Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
                               85                               90                               95

Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
                               100                               105                               110

Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
                               115                               120                               125

Ala Lys Arg Cys Gly Cys Ile
 130                               135

```

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 104 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

```

Met Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile
 1                               5                               10                               15

His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
 20                               25                               30

Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
 35                               40                               45

Tyr Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser
 50                               55                               60

Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
 65                               70                               75                               80

Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
 85                               90                               95

Ser Ala Lys Arg Cys Gly Cys Ile
 100

```

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 103 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

```

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1              5              10              15
Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
      20              25              30
Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
      35              40              45
Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
      50              55              60
Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
      65              70              75              80
Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
      85              90              95
Ala Lys Arg Cys Gly Cys Ile
      100

```

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 114 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

```

Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1              5              10              15
Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
      20              25              30
Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys
      35              40              45
Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys
      50              55              60
Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala
      65              70              75              80
Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser Phe Leu Asp Asp
      85              90              95

```

A-357

- 82 -

Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	Lys	Arg	Cys	Gly
			100					105						110	

Cys Ile

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Hu, Sylvia
- (ii) TITLE OF INVENTION: Truncated Glial Cell Line-Derived
Neurotrophic Factor
- (iii) NUMBER OF SEQUENCES: 50
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: AMGEN INC.
 - (B) STREET: 1840 DeHavilland Drive
 - (C) CITY: Thousand Oaks
 - (D) STATE: California
 - (E) COUNTRY: United States of America
 - (F) ZIP: 91320
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Curry, Daniel R.
 - (B) REGISTRATION NUMBER: 32,727
 - (C) REFERENCE/DOCKET NUMBER: A-357
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 805-447-8102
 - (B) TELEFAX: 805-499-8011
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 402 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 1..402

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCA	CCA	GAT	AAA	CAA	ATG	GCA	GTG	CTT	CCT	AGA	AGA	GAG	CGG	AAT	CGG	48
Ser	Pro	Asp	Lys	Gln	Met	Ala	Val	Leu	Pro	Arg	Arg	Glu	Arg	Asn	Arg	
1				5					10					15		
CAG	GCT	GCA	GCT	GCC	AAC	CCA	GAG	AAT	TCC	AGA	GGA	AAA	GGT	CGG	AGA	96
Gln	Ala	Ala	Ala	Ala	Asn	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	
			20					25					30			
GGC	CAG	AGG	GGC	AAA	AAC	CGG	GGT	TGT	GTC	TTA	ACT	GCA	ATA	CAT	TTA	144
Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	
			35				40					45				
AAT	GTC	ACT	GAC	TTG	GGT	CTG	GGC	TAT	GAA	ACC	AAG	GAG	GAA	CTG	ATT	192
Asn	Val	Thr	Asp	Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	Leu	Ile	
	50					55					60					
TTT	AGG	TAC	TGC	AGC	GGC	TCT	TGC	GAT	GCA	GCT	GAG	ACA	ACG	TAC	GAC	240
Phe	Arg	Tyr	Cys	Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	Tyr	Asp	
65				70					75					80		
AAA	ATA	TTG	AAA	AAC	TTA	TCC	AGA	AAT	AGA	AGG	CTG	GTG	AGT	GAC	AAA	288
Lys	Ile	Leu	Lys	Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	Asp	Lys	
				85					90					95		
GTA	GGG	CAG	GCA	TGT	TGC	AGA	CCC	ATC	GCC	TTT	GAT	GAT	GAC	CTG	TCG	336
Val	Gly	Gln	Ala	Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	Leu	Ser	
			100					105					110			
TTT	TTA	GAT	GAT	AAC	CTG	GTT	TAC	CAT	ATT	CTA	AGA	AAG	CAT	TCC	GCT	384
Phe	Leu	Asp	Asp	Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	
	115					120						125				
AAA	AGG	TGT	GGA	TGT	ATC											402
Lys	Arg	Cys	Gly	Cys	Ile											
	130															

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 134 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ser	Pro	Asp	Lys	Gln	Met	Ala	Val	Leu	Pro	Arg	Arg	Glu	Arg	Asn	Arg	
1				5					10					15		
Gln	Ala	Ala	Ala	Ala	Asn	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	
			20					25					30			
Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	
	35						40					45				

A-357

- 63 -

Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile
50 55 60
Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp
65 70 75 80
Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys
85 90 95
Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser
100 105 110
Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala
115 120 125
Lys Arg Cys Gly Cys Ile
130

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Lys Asn Arg Gly
1

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly
1			5					10	

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Gly	Arg	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly
1			5					10		

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Lys	Gly	Arg	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly
1				5					10		

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly	Lys	Gly	Arg	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly
1				5					10			

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
1 5 10 15

Gly

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
1 5 10 15

Arg Gly

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
1 5 10 15

Asn Arg Gly

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1 5 10 15
 Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
 1 5 10 15
 Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
 1 5 10 15
 Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
1 5 10 15

Gln Arg Gly Lys Asn Arg Gly
20

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg
1 5 10 15

Gly Gln Arg Gly Lys Asn Arg Gly
20

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
1 5 10 15

Arg Gly Gln Arg Gly Lys Asn Arg Gly
20 25

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Asn Arg Gln Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly
 1 5 10 15

Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys
 1 5 10 15

Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly
 1 5 10 15

Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg
 1 5 10 15

Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Asn Pro Glu Asn Ser
 1 5 10 15
 Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn
 1 5 10 15
 Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu
 1 5 10 15

Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro
 1 5 10 15
 Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
 20 25 30
 Gly

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn
 1 5 10 15
 Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
 20 25 30
 Arg Gly

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

```

Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala
1          5          10          15

Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
20          25          30

Asn Arg Gly
35

```

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

```

Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala
1          5          10          15

Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
20          25          30

Lys Asn Arg Gly
35

```

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 37 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

```

Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala
1          5          10          15

Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
20          25          30

Gly Lys Asn Arg Gly
35

```

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

```

Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala
 1             5             10             15
Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
          20             25             30
Arg Gly Lys Asn Arg Gly
          35

```

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

```

Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln
 1             5             10             15
Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
          20             25             30
Gln Arg Gly Lys Asn Arg Gly
          35

```

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 417 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

```

CATATGTCTC CGGATAAACA AATGGCTGTT CTTCCACGTC GTGAACGTAA CCGTCAGGCG 60
GCCGCTGCTA ACCCGGAGAA TTCCCGTGGT AAAGGTCGTC GTGGTCAGCG TGGTAAAAAC 120
CGCGGTTGCG TTCTGACCGC TATCCACCTG AACGTTACCG ACCTGGGTCT CGGTTACGAA 180
ACCAAAGAAG AATTAATCTT CCGTTACTGC TCCGGTTCCT GCGACGCTGC TGAAACCACG 240

```

TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGTCGTC TGGTTTCCGA CAAAGTTGGT 300
 CAAGCTTGCT GCCGTCCGAT CGCTTTCGAC GACGACCTGT CCTTCCTGGA CGACAACCTG 360
 GTTTACCACA TCCTGCGTAA AACTCCCGCT AAGCGTTGCG GTTGCACTTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 417 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CATATGAGCC CGACAAACA GATGCGAGTA CTTCCACGTC GTGAACGTAA TCGCCAGGCA 60
 GCAGCTGCAA ACCCGGAAAA CTCCTGTTGGT AAAGGTCGCC GTGGCCAGCG CGGCAAAAAC 120
 CGTGTTGTG TTCTGACTGC AATCCACCTG AACGTTACTG ACCTGGGTCT GGGCTACGAA 180
 ACCAAAGAAG AACTGATCTT CCGCTACTGC AGCGGCTCTT GCGACGCAGC TGA AACCACT 240
 TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGCCGTC TGTAAGCGA CAAAGTAGGT 300
 CAGGCATGCT GCCGTCCGAT CGCATTCGAC GATGACCTGA GCTTCCTGGA TGACAACCTG 360
 GTTTACCACA TCCTGCGTAA AACTCCCGCT AAACGCTGCG GTTGCACTTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 345 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..342

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ATG	TCC	CCA	GAA	AAT	TCT	CGT	GGT	AAA	GGT	CGT	CGT	GGT	CAG	CGT	GGT	48
Met	Ser	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	Gly	Gln	Arg	Gly	
135					140				145						150	
AAT	AAC	CGC	GGT	TGC	GTT	CTG	ACC	GCT	ATC	CAC	CTG	AAC	GTT	ACC	GAC	96
Asn	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	Asn	Val	Thr	Asp	
			155						160						165	

```

CTG GGT CTC GGT TAC GAA ACC AAA GAA GAA TTA ATC TTC CGT TAC TGC 144
Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys
      170                      175                      180

TCC GGT TCC TGC GAC GCT GCT GAA ACC ACG TAC GAC AAA ATC CTG AAA 192
Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys
      185                      190                      195

AAC CTG TCC CGT AAC CGT CGT CTG GTT TCC GAC AAA GTT GGT CAA GCT 240
Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala
      200                      205                      210

TGC TGC CGT CCG ATC GCT TTC GAC GAC GAC CTG TCC TTC CTG GAC GAC 288
Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp CTG TCC TTC CTG GAC GAC
      215                      220                      225                      230

AAC CTG GTT TAC CAC ATC CTG CGT AAA CAC TCC GCT AAG CGT TGC GGT 336
Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala Lys Arg Cys Gly
      235                      240                      245

TGC ATC TAA
Cys Ile
                                           345

```

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

```

Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1          5          10          15

Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
      20          25          30

Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys
      35          40          45

Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys
      50          55          60

Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala
      65          70          75          80

Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser Phe Leu Asp Asp
      85          90          95

Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala Lys Arg Cys Gly
      100          105          110

Cys Ile

```

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 315 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..312

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

ATG	CGT	GGT	CAA	CGT	GGT	AAA	AAC	CGC	GGT	TGC	GTT	CTG	ACT	GCA	ATC	48
Met	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	
115					120					125					130	
CAC	CTG	AAC	GTT	ACT	GAC	CTG	GGT	CTG	GGC	TAC	GAA	ACC	AAA	GAA	GAA	96
His	Leu	Asn	Val	Thr	Asp	Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	
				135				140						145		
CTG	ATC	TTC	CGC	TAC	TGC	AGC	GGC	TCT	TGC	GAC	GCA	GCT	GAA	ACC	ACT	144
Leu	Ile	Phe	Arg	Tyr	Cys	Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	
			150				155						160			
TAC	GAC	AAA	ATC	CTG	AAA	AAC	CTG	TCC	CGT	AAC	CGC	CGT	CTG	GTA	AGC	192
Tyr	Asp	Lys	Ile	Leu	Lys	Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	
			165				170					175				
GAC	AAA	GTA	GGT	CAG	GCA	TGC	TGC	CGT	CCG	ATC	GCA	TTC	GAC	GAT	GAC	240
Asp	Lys	Val	Gly	Gln	Ala	Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	
	180					185					190					
CTG	AGC	TTC	CTG	GAT	GAC	AAC	CTG	GTT	TAC	CAC	ATC	CTG	CGT	AAA	CAC	288
Leu	Ser	Phe	Leu	Asp	Asp	Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	
195				200					205					210		
TCC	GCT	AAA	CGC	TGC	GGT	TGC	ATC	TAA								315
Ser	Ala	Lys	Arg	Cys	Gly	Cys	Ile									
				215												

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Met	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile
1					5				10					15	

- 78 -

(2) INFORMATION FOR SEQ ID NO:45:

(A) LENGTH: 312 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..309

ATG	GGT	CAA	CGT	GGT	AAA	AAC	CGT	GGT	TGT	GTT	CTG	ACT	GCA	ATC	CAC	48
Met	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	
105					110					115				120		
CTG	AAC	GTT	ACT	GAC	CTG	GGT	CTG	GGC	TAC	GAA	ACC	AAA	GAA	GAA	CTG	96
Leu	Asn	Val	Thr	Asp	Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	Leu	
				125					130					135		
ATC	TTC	CGC	TAC	TGC	AGC	GGC	TCT	TGC	GAC	GCA	GCT	GAA	ACC	ACT	TAC	144
Ile	Phe	Arg	Tyr	Cys	Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	Tyr	
			140					145					150			
GAC	AAA	ATC	CTG	AAA	AAC	CTG	TCC	CGT	AAC	CGC	CGT	CTG	GTA	AGC	GAC	192
Asp	Lys	Ile	Leu	Lys	Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	Asp	
		155					160					165				
AAA	GTA	GGT	CAG	GCA	TGC	TGC	CGT	CCG	ATC	GCA	TTC	GAC	GAT	GAC	CTG	240
Lys	Val	Gly	Gln	Ala	Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	Leu	
	170				175						180					
AGC	TTC	CTG	GAT	GAC	AAC	CTG	GTT	TAC	CAC	ATC	CTG	CGT	AAA	CAC	TCC	288
Ser	Phe	Leu	Asp	Asp	Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	
185					190					195				200		

GCT AAA CGC TGC GGT TGC ATC TAA
 Ala Lys Arg Cys Gly Cys Ile
 205

312

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1 5 10 15
 Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 20 25 30
 Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 35 40 45
 Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
 50 55 60
 Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
 65 70 75 80
 Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
 85 90 95
 Ala Lys Arg Cys Gly Cys Ile
 100

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 135 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Met Ser Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn
 1 5 10 15
 Arg Gln Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
 20 25 30
 Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 35 40 45

```

Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 50                      55                      60

Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
65                      70                      75                      80

Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
                      85                      90                      95

Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
100                      105                      110

Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
115                      120                      125

Ala Lys Arg Cys Gly Cys Ile
130                      135

```

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

```

Met Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile
 1                      5                      10                      15

His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
20                      25                      30

Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
35                      40                      45

Tyr Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser
50                      55                      60

Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
65                      70                      75                      80

Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
85                      90                      95

Ser Ala Lys Arg Cys Gly Cys Ile
100

```

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 103 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

```

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1           5           10           15
Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Leu
          20           25           30
Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
          35           40           45
Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
          50           55           60
Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
65           70           75           80
Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
          85           90           95
Ala Lys Arg Cys Gly Cys Ile
          100

```

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

```

Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1           5           10           15
Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
          20           25           30
Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys
          35           40           45
Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys
          50           55           60
Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala
65           70           75           80
Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser Phe Leu Asp Asp
          85           90           95

```

A-357

- 82 -

Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	Lys	Arg	Cys	Gly
			100					105						110	

Cys Ile